




St. Vrain Valley School District High School

2009 **Youth Risk Behavior Survey (YRBS)**

PARTNERS



Boulder Valley School District
6500 Arapahoe
Boulder, CO 80303



St. Vrain Valley School District
395 South Pratt Parkway
Longmont, CO 80501



Health Planning Program
Boulder County Public Health
3450 Broadway
Boulder, CO 80304



Healthy Youth Alliance of Boulder County
Boulder County Community Services
3481 Broadway
Boulder, CO 80304



FOREWORD

In late 2009, Boulder Valley School District (BVSD), St. Vrain Valley School District (SVVSD), Boulder County Healthy Youth Alliance (HYA), and Boulder County Public Health (BCPH) advanced our collaborative effort to learn more about how we can protect and enhance the well-being of our county's school-age youth. Together, we conducted the fifth Youth Risk Behavior Survey (YRBS) among a representative sample of Boulder County high school students. Additionally, in BVSD, a middle school survey was administered to 7th and 8th grade students to further improve our ability to promote healthy decision-making and prevent behaviors that can lead to long-lasting, negative impacts.

The data presented in this report reflect a single point in time when the survey was conducted; however, comparing this data to previous YRBS data, we can begin to identify more long-term trends that help in planning interventions. Since the original survey was conducted in 2001, the data have been used to develop school- and community-based strategies that have been implemented to reduce behavioral risk factors along adolescents. Whether strengthening the health curriculum or addressing disparities among racial, ethnic, or sexual orientation lines, we have seen many improvements and yet still face many challenges. Continued efforts that strengthen the knowledge and skills of our children, as well as that of our youths' parents and other community members, can help promote preventive behavior and reduce risk behaviors among our youth.

As we look at these data, four preliminary insights emerge. These insights can be useful as a guide to understanding and responding to these data:

- Parental influence can help reduce or even prevent many risk behaviors (e.g., smoking, alcohol use, and marijuana use).
- There are many positive values, beliefs, attitudes, and behaviors (social norms) that we can reinforce and promote (e.g., use of seatbelts).
- There are many risk behaviors that, with effective interventions, can be prevented, reduced, or postponed (e.g., alcohol use, binge drinking, and driving or riding with a driver who has consumed alcohol).
- There are some risk factors that call for immediate intervention and help (e.g., forced sexual intercourse, clinical depression, and suicide ideation).

This information provides the school districts with a valuable opportunity to partner with the communities in which our youth reside and to support our youth as they move toward the future as healthy, educated, and productive adults.

For additional information, please visit www.BoulderCountyYRBS.org.

Robin Bohannon
Director
Boulder County
Community Services

Don Haddad
Superintendent
St. Vrain Valley
School District

Bruce Messinger
Superintendent
Boulder Valley
School District

Jeffrey Zayach
Director
Boulder County
Public Health

Introduction

Data from the Youth Risk Behavior Survey (YRBS) is used to examine the risk and preventative behaviors among adolescents in middle and high schools across the country as an indicator of health. In addition, the data is used as an evaluative tool to monitor the progress of school- and community-based strategies that are being implemented to improve and/or maintain the health of our youth. The data are a catalyst for the community process that has emerged to address the health and well-being of our youth. Although the data tell us what the youth are doing, it is the community response that creates change and positively impacts our youth.

The 2009 Boulder County YRBS follows YRBS surveys conducted in Boulder County high schools in both the Boulder Valley (BVSD) and St. Vrain Valley (SVVSD) School Districts in 2003, 2005, and 2007. The 2009 YRBS survey marked the first time that data were collected at the middle school level in Boulder County. These data were collected from middle schools in BVSD and included students in the 7th and 8th grades. Data from the 2009 YRBS are presented in four reports: one each for St. Vrain Valley School District and Boulder Valley School District high schools, one for Boulder Valley School District middle schools, and a Boulder County report that combines the results from the high schools in each of the two school districts.

2009 YRBS results

All reports, background information, and data from the 2009 Boulder County YRBS for each school district, as well as data for Boulder County high schools overall, can be found at www.BoulderCountyYRBS.org.

The results are generalizable to the schools from which each sample was selected. Data are presented by grade, gender, race, and ethnicity (i.e., “Hispanic,” “white non-Hispanic,” or “other non-Hispanic”). Since 2007, data collected on ethnicity have been separated from race, allowing analysis to be conducted specifically for Hispanic youth, increasing our ability to identify health and behavior disparities. Data about sexual orientation (i.e., “heterosexual” or “LGBQ” [lesbian, gay, bisexual, or questioning]) are only reported at the county level to protect the confidentiality of the LGBQ students. The results of these stratified analyses provide an opportunity to examine and address disparities in health risk behaviors. They also reinforce the importance of assuring that community efforts address the needs of ALL Boulder County youth.

Data from the 2009 Colorado YRBS and the 2009 national YRBS are included in appendices where available. While every effort was made to collect and analyze the data using similar protocols, these data are not directly comparable and should be used for reference only. Colorado YRBS data and methods are detailed at <http://hkcs.omni.org/state>; national YRBS data and methods are described at www.cdc.gov/HealthyYouth/yrbs/index.htm.

Conducting the YRBS

The Boulder County YRBS survey procedures are based on the Centers for Disease Control and Prevention (CDC) protocol (www.cdc.gov/HealthyYouth/yrbs/yrbs_methods.htm).

Sample selection

Nine SVVSD high schools, 9 BVSD high schools, and 16 BVSD middle schools participated in the 2009 YRBS. Each school chose a day and class period in early November 2009 for survey administration. Classes from each of these periods were randomly selected to be surveyed. This selection yielded 66 SVVSD high school and 67 BVSD high school classrooms. A separate sample selection identified 41 BVSD middle school classrooms in which 7th and 8th grade students would have the opportunity to take the middle school survey. Information was mailed to the parents of each student in the selected classrooms to inform them about the survey, student selection, and how they could ask questions and/or exclude their students from participating if they wished.

Survey administration

Data for the 2009 survey were collected during the week of November 9, 2009, for BVSD middle schools, and the week of November 16, 2009, for high schools in both BVSD and SVVSD. Survey implementation procedures were designed to protect students' privacy by allowing for anonymous and voluntary participation. Participating students completed the survey during regular class time, under the supervision of trained volunteers.

Data analysis

Prevalence estimates and 95% confidence intervals were calculated and are presented in the data tables. The high response rate enabled data to be weighted by gender and grade level so that the results are representative of the entire population in schools from which each sample was selected. Therefore, the percentages reported are estimates of the occurrence of the behaviors in the *entire population* from which the sample was selected, not just for the students *who took the survey*.

Data tables

A sample data table appears on the next page. Percentages presented in the tables are *prevalence estimates*. Prevalence estimates have been replaced by "***" where numbers were too small to report for statistical or confidentiality reasons and "---" when data analysis was not completed for the specified category for confidentiality reasons or because the question was not asked. Also included are *95% confidence intervals* (in parentheses below the prevalence estimates). These intervals indicate the range in which the *actual prevalence* (when generalized from the sample to the survey population) was likely to fall. The size of the confidence interval reflects the certainty of the accuracy of the estimate. Larger confidence intervals reflect less certainty; smaller confidence intervals reflect greater certainty. Confidence intervals may be used to determine the statistical significance of differences in observed results. Confidence intervals that do not overlap indicate that the difference in estimates is statistically significant (assuming a significance level of $\alpha = 0.05$).

For more information

www.BoulderCountyYRBS.org



ST. VRAIN VALLEY SCHOOL DISTRICT SAMPLE DATA TABLE

Confidence interval

Prevalence estimate for all SVVSD 9th graders

Unintentional Injuries - Helmets and Seatbelts

	<i>Never or rarely wore bicycle helmet*</i>			<i>Never or rarely wore seat belt**</i>		
Grade	Total	Female	Male	Total	Female	Male
9th	80.7% (78.4-82.8)	79.2% (75.5-82.5)	82.1% (79.2-84.7)	7.6% (6.3-9.2)	6.9% (5.0-9.4)	8.3% (6.6-10.5)
10th	75.7% (73.3-78.0)	73.5% (69.8-77.0)	77.8% (74.4-80.9)	5.9% (4.6-7.6)	3.5% (2.3-5.5)	8.2% (6.0-11.0)
11th	78.3% (76.0-80.4)	79.5% (76.1-82.5)	77.5% (74.2-80.4)	6.0% (4.5-8.0)	3.8% (1.9-7.6)	8.1% (6.2-10.5)
12th	74.8% (71.2-78.0)	76.1% (71.0-80.5)	73.9% (69.0-78.3)	4.8% (3.6-6.5)	3.4% (2.0-5.8)	6.3% (4.4-8.8)
Race/Ethnicity						
White non-Hispanic	71.5% (69.8-73.1)	---	---	3.8% (3.2-4.5)	---	---
Hispanic	92.3% (90.1-94.0)	---	---	11.9% (9.8-14.4)	---	---
Other non-Hispanic	83.5% (80.1-86.4)	---	---	5.6% (3.5-8.9)	---	---
Total	77.6% (76.3-78.8)	77.1% (75.2-78.9)	78.0% (76.2-79.7)	6.1% (5.4-6.9)	4.4% (3.5-5.6)	7.7% (6.7-8.9)

* Among the 83.3% of students who rode a bicycle during the 12 months preceding the survey
 ** When riding in a car driven by someone else in the 30 days preceding the survey
 --- Not available

Prevalence estimate for all SVVSD students

Prevalence estimate for all SVVSD female students

Prevalence estimate for all SVVSD male students



St. Vrain Valley School District (SVVSD) High School Narrative

The narrative following each data table in this report consists of one paragraph for each survey question. This paragraph reports the total prevalence estimate for each behavior, followed by all statistically significant differences and only statistically significant differences by race/ethnicity total, sexual orientation total, and gender total, as well as grade if it changed significantly in the same direction (increase or decrease) from 9th to 10th grade, 10th to 11th grade, and 11th to 12th grade. If differences in the totals are not noted in the narrative, prevalence estimates are not statistically different and should not be interpreted or reported as such. If there are no statistically significant differences in the totals, the narrative includes only the total prevalence estimate. Statistically significant differences among the gender and grade strata subtotals (e.g., 9th and 10th grade or 9th grade females and 9th grade males) may exist but are not reported in the narrative.

St. Vrain Valley School District (SVVSD) High School Results

There were 1,636 high school students in 66 classrooms sampled to participate in the 2009 St. Vrain Valley School District YRBS. Completed surveys were collected from 1,221 SVVSD high school students and 9 schools participated, yielding an overall response rate of 75%. The following table summarizes the weighted demographics of the respondents.

Sex		Grade		Race/Ethnicity	
Female	49.0%	9 th	25.5%	White, non-Hispanic	65.2%
Male	51.0%	10 th	24.4%	Hispanic	24.7%
		11 th	25.6%	Other, non-Hispanic	10.2%
		12 th	24.6%		

The overall response rate of 75% allowed for weighting of the data to represent all students in grades 9-12 at the participating high schools. In essence, this means that the prevalence rates presented in this report are estimates of the prevalence in this entire population of high school students; they are not merely the prevalence rates within the sample group of students.



Family and Community Service

	<i>Parents have clear rules and standards for behavior</i>			<i>Ate 4 or more meals with family*</i>			<i>Did community service **</i>		
Grade	Total	Female	Male	Total	Female	Male	Total	Female	Male
9th	80.6% (78.3-82.8)	78.1% (74.5-81.3)	83.2% (80.1-85.8)	79.1% (76.7-81.3)	76.0% (72.5-79.1)	82.1% (78.7-85.1)	37.4% (34.7-40.1)	43.7% (39.8-47.6)	31.2% (27.9-34.7)
10th	82.8% (80.6-84.9)	85.5% (82.7-88.0)	80.2% (76.7-83.4)	80.2% (77.9-82.4)	78.1% (75.1-80.9)	82.3% (78.7-85.4)	39.1% (36.5-41.7)	41.0% (37.3-44.9)	37.2% (33.5-40.9)
11th	85.6% (83.6-87.5)	84.0% (80.9-86.6)	87.2% (84.2-89.6)	64.2% (61.4-66.9)	61.6% (57.5-65.5)	66.6% (62.7-70.2)	45.9% (43.2-48.7)	50.6% (46.5-54.7)	41.7% (38.0-45.5)
12th	89.6% (86.8-91.8)	94.9% (91.3-97.1)	84.3% (79.7-88.0)	69.4% (66.1-72.5)	64.0% (59.0-68.7)	74.7% (70.4-78.5)	45.3% (42.0-48.7)	55.7% (50.6-60.6)	35.3% (31.0-39.8)
Race/Ethnicity									
White non-Hispanic	86.6% (85.2-87.9)	---	---	77.5% (75.9-79.0)	---	---	45.7% (43.9-47.5)	---	---
Hispanic	80.4% (77.9-82.7)	---	---	63.7% (60.6-66.6)	---	---	31.4% (28.6-34.4)	---	---
Other non-Hispanic	83.7% (79.9-87.0)	---	---	69.2% (64.8-73.3)	---	---	43.2% (38.9-47.7)	---	---
Total	84.7% (83.5-85.7)	85.6% (84.0-87.0)	83.8% (82.1-85.3)	73.2% (71.8-74.5)	69.9% (68.0-71.8)	76.3% (74.4-78.0)	41.9% (40.5-43.4)	47.7% (45.6-49.9)	36.4% (34.5-38.3)

* During the 7 days preceding the survey

** In the 30 days preceding the survey

--- Not available

More than 4 in 5 students in surveyed SVVSD high schools (84.7%) reported they believed their parents or other adults in their family had clear rules and standards for their behavior. Overall, prevalence of this belief was higher among white non-Hispanic students (86.6%) than it was among Hispanic students (80.4%).

More than 7 in 10 students in surveyed SVVSD high schools (73.2%) reported they had eaten 4 or more meals with their families during the 7 days prior to the survey. Overall, prevalence of having eaten 4 or more meals with their families was higher among white non-Hispanic students (77.5%) than both other non-Hispanic and Hispanic students (69.2% and 63.7%, respectively); it was also higher among male students (76.3%) than it was among female students (69.9%).

More than 4 in 10 students in surveyed SVVSD high schools (41.9%) reported they had performed organized community service as a non-paid volunteer during the 30 days prior to the survey. Overall, prevalence for having performed some kind of organized community service as a non-paid volunteer was higher among both white non-Hispanic and other non-Hispanic students (45.7% and 43.2% respectively) than Hispanic students (31.4%); it was also higher among female students (47.7%) than male students (36.4%).



Unintentional Injuries - Helmets and Seatbelts

Grade	Never or rarely wore bicycle helmet*			Never or rarely wore seat belt**		
	Total	Female	Male	Total	Female	Male
9th	80.7% (78.4-82.8)	79.2% (75.5-82.5)	82.1% (79.2-84.7)	7.6% (6.3-9.2)	6.9% (5.0-9.4)	8.3% (6.6-10.5)
10th	75.7% (73.3-78.0)	73.5% (69.8-77.0)	77.8% (74.4-80.9)	5.9% (4.6-7.6)	3.5% (2.3-5.5)	8.2% (6.0-11.0)
11th	78.3% (76.0-80.4)	79.5% (76.1-82.5)	77.5% (74.2-80.4)	6.0% (4.5-8.0)	3.8% (1.9-7.6)	8.1% (6.2-10.5)
12th	74.8% (71.2-78.0)	76.1% (71.0-80.5)	73.9% (69.0-78.3)	4.8% (3.6-6.5)	3.4% (2.0-5.8)	6.3% (4.4-8.8)
Race/Ethnicity						
White non-Hispanic	71.5% (69.8-73.1)	---	---	3.8% (3.2-4.5)	---	---
Hispanic	92.3% (90.1-94.0)	---	---	11.9% (9.8-14.4)	---	---
Other non-Hispanic	83.5% (80.1-86.4)	---	---	5.6% (3.5-8.9)	---	---
Total	77.6% (76.3-78.8)	77.1% (75.2-78.9)	78.0% (76.2-79.7)	6.1% (5.4-6.9)	4.4% (3.5-5.6)	7.7% (6.7-8.9)

* Among the 79.2% of students who rode a bicycle during the 12 months preceding the survey

** When riding in a car driven by someone else in the 30 days preceding the survey

--- Not available

Among students in surveyed SVVSD high schools who reported they had ridden a bicycle in the 12 months prior to the survey, 77.6% reported they had rarely or never wore a helmet. Overall, the prevalence of never or rarely wearing a bicycle helmet was higher among Hispanic students (92.3%) than other non-Hispanic students (83.5%), and both were higher than white non-Hispanic students (71.5%).

Overall, 6.1% of students in surveyed SVVSD high schools reported they rarely or never wore a seatbelt when they rode in a car driven by someone else. Overall, prevalence of this behavior was higher among Hispanic students (11.9%) than both other non-Hispanic (5.6%) and white non-Hispanic students (3.8%); it was also higher among male students (7.7%) than female students (4.4%).

Drinking and Driving

	<i>Rode with a driver who had been drinking alcohol*</i>			<i>Drove after drinking alcohol*</i>		
Grade	Total	Female	Male	Total	Female	Male
9th	25.9% (23.6-28.5)	31.1% (27.4-35.1)	20.9% (18.1-24.0)	5.4% (4.1-7.0)	6.8% (4.9-9.3)	4.0% (2.5-6.2)
10th	21.3% (19.0-23.8)	26.5% (23.1-30.3)	16.1% (13.3-19.4)	4.7% (3.7-6.1)	3.5% (2.2-5.6)	5.9% (4.3-7.9)
11th	22.7% (20.4-25.2)	21.0% (17.8-24.5)	24.3% (21.1-27.9)	8.5% (7.0-10.3)	4.9% (3.5-6.8)	11.8% (9.4-14.7)
12th	24.1% (21.2-27.3)	19.8% (16.0-24.2)	28.5% (24.1-33.3)	13.9% (11.5-16.7)	11.9% (8.6-16.2)	15.8% (12.4-20.1)
Race/Ethnicity						
White non-Hispanic	21.0% (19.5-22.6)	---	---	7.3% (6.3-8.5)	---	---
Hispanic	32.4% (29.5-35.3)	---	---	11.0% (9.2-13.2)	---	---
Other non-Hispanic	19.1% (15.6-23.1)	---	---	6.0% (4.2-8.6)	---	---
Total	23.5% (22.3-24.9)	24.6% (22.8-26.5)	22.5% (20.8-24.3)	8.1% (7.3-9.0)	6.8% (5.7-8.2)	9.4% (8.1-10.8)

* One or more times during the 30 days preceding the survey

--- Not available

More than 1 in 5 students in surveyed SVVSD high schools (23.5%) reported that, in the 30 days preceding the survey, they had ridden in a car or other vehicle driven by someone who had been drinking alcohol. Overall, prevalence of having ridden in a car or other vehicle driven by someone who had been drinking alcohol was higher among Hispanic students (32.4%) than both white non-Hispanic (21.0%) and other non-Hispanic students (19.1%).

Among students in surveyed SVVSD high schools, 8.1% reported that, in the 30 days before the survey, they had driven a car or other vehicle after they had been drinking alcohol. Overall, prevalence of having driven a vehicle after drinking alcohol was higher among Hispanic students (11.0%) than both white non-Hispanic (7.3%) and other non-Hispanic students (6.0%).

Safety

	<i>Felt safe in their neighborhood</i>			<i>Did not go to school because felt unsafe*</i>		
Grade	Total	Female	Male	Total	Female	Male
9th	84.7% (82.4-86.8)	82.9% (79.1-86.0)	86.6% (83.7-89.0)	6.4% (5.0-8.0)	10.9% (8.4-14.0)	2.0% (1.3-3.1)
10th	86.0% (83.8-88.0)	87.4% (84.2-90.0)	84.7% (81.3-87.5)	5.2% (4.0-6.8)	7.7% (5.5-10.6)	2.8% (1.8-4.4)
11th	88.4% (86.2-90.3)	90.3% (86.8-93.0)	86.7% (83.7-89.2)	5.0% (3.9-6.4)	6.4% (4.6-8.7)	3.7% (2.6-5.4)
12th	93.5% (91.3-95.2)	95.6% (91.7-97.7)	91.4% (88.5-93.7)	3.9% (2.5-6.1)	2.9% (1.2-7.2)	4.8% (3.1-7.6)
Race/Ethnicity						
White non-Hispanic	91.3% (90.1-92.4)	---	---	3.8% (3.0-4.8)	---	---
Hispanic	82.1% (79.5-84.5)	---	---	10.0% (8.4-12.0)	---	---
Other non-Hispanic	82.9% (78.8-86.4)	---	---	2.3% (1.3-4.3)	---	---
Total	88.2% (87.1-89.2)	89.0% (87.4-90.5)	87.3% (85.9-88.6)	5.1% (4.4-5.9)	7.0% (5.8-8.4)	3.3% (2.7-4.2)

* At school (or on the way to/from school) during 30 days preceding the survey

--- Not available

Nearly 9 in 10 students in surveyed SVVSD high schools (88.2%) reported feeling safe in their neighborhood always or most of the time. Overall, the prevalence of feeling safe in their neighborhood always or most of the time was higher among white non-Hispanic students (91.3%) than both other non-Hispanic (82.9%) and Hispanic students (82.1%).

More than 1 in 20 students in surveyed SVVSD high schools (5.1%) reported that, during the 30 days prior to the survey, they had not gone to school because they felt they would be unsafe at school or on the way to or from school. Overall, the prevalence of not going to school because they felt they would be unsafe at school or on the way to or from school was higher among Hispanic students (10.0%) than both white non-Hispanic and other non-Hispanic students (3.8% and 2.3%, respectively); it was also higher among female students than male students (7.0% and 3.3%, respectively).



Harassment at School (or on the way to/from school) – Race/Ethnicity or Sexual Orientation

Grade	<i>Harassed*</i>			<i>Harassed because of race or ethnic origin*</i>			<i>Harassed because someone thought you were lesbian, gay, or bisexual*</i>		
	Total	Female	Male	Total	Female	Male	Total	Female	Male
9th	35.9% (33.2-38.6)	38.9% (34.9-43.0)	32.9% (29.4-36.6)	7.4% (5.9-9.1)	6.8% (4.7-9.7)	7.9% (6.2-10.0)	9.5% (7.9-11.4)	9.1% (6.7-12.3)	9.8% (7.8-12.2)
10th	28.4% (25.9-31.2)	30.7% (26.9-34.8)	26.5% (23.0-30.3)	7.5% (5.9-9.3)	5.7% (3.9-8.4)	9.1% (6.8-12.2)	8.6% (7.0-10.4)	6.9% (4.9-9.6)	10.2% (8.0-12.9)
11th	28.6% (26.0-31.3)	31.4% (27.7-35.5)	26.0% (22.5-29.7)	5.8% (4.5-7.4)	4.4% (2.9-6.5)	7.0% (5.1-9.7)	7.5% (6.2-9.1)	7.1% (5.3-9.5)	7.9% (6.0-10.2)
12th	20.0% (17.2-23.2)	16.8% (13.1-21.2)	22.9% (19.0-27.5)	4.3% (2.9-6.4)	*** ***	6.8% (4.3-10.6)	4.5% (3.1-6.5)	*** ***	8.0% (5.3-11.7)
Race/Ethnicity									
White non-Hispanic	29.4% (27.7-31.2)	---	---	3.0% (2.3-3.8)	---	---	7.5% (6.6-8.6)	---	---
Hispanic	25.7% (23.0-28.6)	---	---	12.9% (10.9-15.2)	---	---	7.7% (6.3-9.5)	---	---
Other non-Hispanic	26.7% (22.7-31.0)	---	---	11.7% (9.0-15.0)	---	---	7.5% (5.4-10.3)	---	---
Total	28.2% (26.9-29.6)	29.6% (27.6-31.6)	27.0% (25.2-28.9)	6.2% (5.5-7.1)	4.7% (3.8-5.8)	7.7% (6.5-9.1)	7.5% (6.8-8.4)	6.0% (5.1-7.2)	8.9% (7.8-10.3)

* At school (or on the way to/from school) during the 12 months preceding the survey

*** Numerator <3 and/or denominator <30

--- Not available

Harassment can include threatening, bullying, name calling or obscenities, offensive notes or graffiti, exclusion from groups, unwanted attention, or unwanted touching and physical assault. During the 12 months preceding the survey, more than 1 in 4 students in surveyed SVVSD high schools (28.2%) reported they had been harassed at school (or on the way to or from school).

In the 12 months prior to the survey, more than 1 in 20 students in surveyed SVVSD high schools (6.2%) reported they had been harassed because of their race or ethnic origin at school (or on the way to or from school). Overall, the prevalence of harassment because of their race or ethnic origin at school was higher among both Hispanic (12.9%) and other non-Hispanic students (11.7%) than white non-Hispanic students (3.0%); it was also higher among male students than female students (7.7% and 4.7%, respectively).

During the 12 months prior to the survey, 7.5% of students in surveyed SVVSD high schools reported they had been harassed at school (or on the way to or from school) because someone thought they were gay, lesbian, or bisexual; it was also higher among male students (8.9%) than female students (6.0%).



Harassment at School (or on the way to/from school) – Religion, Physical Appearance, Unwanted Sexual Attention

Grade	<i>Harassed because of religious beliefs*</i>			<i>Harassed because of weight, size, physical appearance*</i>			<i>Received unwanted sexual comments or attention*</i>		
	Total	Female	Male	Total	Female	Male	Total	Female	Male
9th	4.3% (3.3-5.5)	4.7% (3.3-6.8)	3.8% (2.8-5.3)	16.0% (14.0-18.3)	18.7% (15.5-22.3)	13.5% (11.1-16.3)	21.5% (19.3-23.8)	35.3% (31.7-39.1)	8.0% (6.2-10.2)
10th	4.6% (3.5-5.9)	5.6% (4.0-7.8)	3.5% (2.3-5.3)	13.4% (11.4-15.7)	15.0% (12.0-18.5)	12.0% (9.5-15.1)	19.4% (17.3-21.7)	34.0% (30.4-37.8)	5.2% (3.7-7.2)
11th	4.3% (3.4-5.5)	4.9% (3.5-6.6)	3.9% (2.7-5.6)	12.8% (11.0-14.8)	13.6% (11.0-16.6)	12.0% (9.8-14.7)	16.6% (14.6-18.8)	27.9% (24.4-31.8)	6.3% (4.7-8.3)
12th	4.6% (3.2-6.7)	5.5% (3.2-9.5)	3.8% (2.4-5.8)	10.0% (7.9-12.5)	6.5% (3.9-10.5)	13.3% (10.4-16.8)	17.9% (15.4-20.7)	26.0% (21.9-30.7)	10.1% (7.7-13.2)
Race/Ethnicity									
White non-Hispanic	4.3% (3.5-5.2)	---	---	13.8% (12.5-15.2)	---	---	20.0% (18.6-21.6)	---	---
Hispanic	4.5% (3.5-5.9)	---	---	12.6% (10.6-14.9)	---	---	17.6% (15.5-20.0)	---	---
Other non-Hispanic	5.6% (4.0-7.7)	---	---	8.3% (6.2-11.2)	---	---	14.8% (11.9-18.2)	---	---
Total	4.4% (3.9-5.1)	5.2% (4.2-6.3)	3.8% (3.1-4.6)	13.0% (12.0-14.1)	13.4% (11.9-15.1)	12.7% (11.4-14.1)	18.9% (17.7-20.0)	30.9% (29.0-32.8)	7.4% (6.4-8.5)

* At school (or on the way to/from school) during the 12 months preceding the survey

--- Not available

During the 12 months preceding the survey, 4.4% of students in surveyed SVVSD high schools reported they had been harassed at school (or on the way to or from school) due to religious beliefs.

More than 1 in 10 students in surveyed SVVSD high schools (13.0%) reported that, during the 12 months prior to the survey, they had been harassed at school (or on the way to or from school) because of their weight, size, or physical appearance. Overall, the prevalence of harassment because of weight, size, or physical appearance was higher among white non-Hispanic students (13.8%) than other non-Hispanic students (8.3%).

During the 12 months prior to the survey, 18.9% of students in surveyed SVVSD high schools reported they had received unwanted sexual comments or attention while at school (or on the way to or from school). Overall, the prevalence of receiving unwanted sexual comments or attention was higher among white non-Hispanic students (20.0%) than other non-Hispanic students (14.8%); it was also higher among female students than male students (30.9% and 7.4%, respectively). Additionally, at each of the four grade levels, female students had a higher prevalence of receiving unwanted sexual comments or attention than male students.



Violence Anywhere

Grade	Carried a weapon*			In physical fight**			Injured in fight and needed medical treatment**		
	Total	Female	Male	Total	Female	Male	Total	Female	Male
9th	17.1% (15.2-19.2)	9.7% (7.7-12.2)	24.2% (21.2-27.5)	35.8% (33.3-38.4)	28.7% (25.2-32.4)	42.8% (39.2-46.5)	2.2% (1.5-3.2)	3.2% (2.1-5.0)	***
10th	15.2% (13.3-17.3)	5.9% (4.3-8.1)	24.2% (21.0-27.7)	26.3% (23.9-28.9)	19.2% (16.2-22.5)	33.3% (29.6-37.2)	2.9% (2.0-4.3)	3.1% (1.7-5.5)	2.7% (1.6-4.5)
11th	17.2% (15.2-19.5)	5.0% (3.5-7.0)	28.5% (25.1-32.3)	28.5% (26.0-31.1)	19.1% (16.0-22.7)	37.3% (33.5-41.1)	3.2% (2.2-4.5)	3.2% (1.8-5.6)	3.2% (2.0-4.9)
12th	13.6% (11.2-16.3)	5.5% (2.9-10.2)	21.5% (18.1-25.3)	22.4% (19.8-25.3)	10.1% (7.5-13.5)	34.5% (30.2-39.2)	1.2% (0.7-2.1)	***	2.3% (1.3-4.1)
Race/Ethnicity									
White non-Hispanic	17.5% (16.2-18.9)	---	---	26.9% (25.3-28.5)	---	---	1.9% (1.4-2.5)	---	---
Hispanic	13.5% (11.6-15.8)	---	---	35.4% (32.6-38.3)	---	---	4.0% (2.9-5.5)	---	---
Other non-Hispanic	10.7% (7.8-14.4)	---	---	21.7% (18.2-25.7)	---	---	***	---	---
Total	15.8% (14.7-16.9)	6.5% (5.4-7.9)	24.6% (23.0-26.4)	28.3% (27.1-29.6)	19.3% (17.7-21.1)	37.0% (35.1-39.0)	2.4% (1.9-2.9)	2.4% (1.8-3.2)	2.4% (1.8-3.1)

* On 1 or more of the 30 days preceding the survey
 ** One or more times during the 12 months preceding the survey
 *** Numerator <3 and/or denominator <30
 --- Not available

Nearly 1 in 7 students in surveyed SVVSD high schools (15.8%) reported they had carried a weapon (such as a gun, knife, or club) in the 30 days preceding the survey. Overall, the prevalence of carrying such a weapon was higher among white non-Hispanic students (17.5%) than both Hispanic and other non-Hispanic students (13.5% and 10.7%, respectively); it was also higher among male students (24.6%) than female students (6.5%). Additionally, at each of the four grade levels, male students had a higher prevalence than female students.

More than 1 in 4 students in surveyed SVVSD high schools (28.3%) reported they had been in a physical fight during the 12 months preceding the survey. Overall, the prevalence for having been in a physical fight was higher among Hispanic students (35.4%) than both white non-Hispanic and other non-Hispanic students (26.9% and 21.7%, respectively); it was also higher among male students (37.0%) than female students (19.3%). Additionally, at each of the four grade levels, male students had a higher prevalence for having been in a physical fight than female students.

Not all physical fights result in injuries serious enough to be treated by a doctor or nurse; however, 2.4% of students in surveyed SVVSD high schools reported that, in the 12 months prior to the survey, they had been in a physical fight in which they were injured and had to be treated by a professional. Overall, prevalence for having been in a physical fight that required medical treatment was higher among Hispanic students than white non-Hispanic students (4.0% and 1.9%, respectively).



Violence on School Property

	<i>Carried a weapon on school property*</i>			<i>Threatened or injured with a weapon on school property**</i>			<i>In physical fight on school property**</i>		
Grade	Total	Female	Male	Total	Female	Male	Total	Female	Male
9th	4.3% (3.3-5.5)	4.7% (3.3-6.7)	3.9% (2.7-5.6)	8.4% (6.9-10.2)	7.2% (5.1-10.1)	9.6% (7.4-12.3)	14.2% (12.5-16.1)	11.8% (9.8-14.2)	16.4% (13.8-19.4)
10th	4.2% (3.1-5.7)	2.5% (1.5-3.9)	5.9% (4.0-8.7)	5.9% (4.6-7.6)	4.0% (2.8-5.9)	7.7% (5.5-10.6)	10.2% (8.4-12.2)	8.9% (6.7-11.8)	11.4% (8.8-14.5)
11th	5.5% (4.3-6.9)	2.3% (1.3-4.0)	8.4% (6.5-10.9)	6.6% (5.3-8.3)	3.9% (2.7-5.7)	9.1% (7.0-11.9)	10.8% (9.1-12.8)	6.2% (4.4-8.7)	15.1% (12.4-18.2)
12th	6.3% (4.9-8.1)	*** ***	10.5% (8.2-13.4)	4.5% (3.5-5.7)	*** ***	7.5% (5.7-9.9)	7.4% (5.8-9.3)	*** ***	14.1% (11.2-17.7)
Race/Ethnicity									
White non-Hispanic	5.3% (4.5-6.2)	--- ---	--- ---	6.3% (5.4-7.2)	--- ---	--- ---	8.9% (8.0-10.0)	--- ---	--- ---
Hispanic	5.6% (4.3-7.2)	--- ---	--- ---	7.2% (5.7-9.2)	--- ---	--- ---	15.5% (13.5-17.9)	--- ---	--- ---
Other non-Hispanic	2.6% (1.4-4.8)	--- ---	--- ---	5.3% (3.6-7.7)	--- ---	--- ---	10.6% (8.0-14.0)	--- ---	--- ---
Total	5.1% (4.4-5.8)	2.9% (2.2-3.7)	7.2% (6.2-8.3)	6.4% (5.7-7.1)	4.2% (3.4-5.1)	8.5% (7.4-9.8)	10.7% (9.8-11.6)	6.9% (6.0-8.0)	14.3% (12.9-15.8)

* On 1 or more of the 30 days preceding the survey

** One or more times during the 12 months preceding the survey

*** Numerator <3 and/or denominator <30

--- Not available

More than 1 in 20 students in surveyed SVVSD high schools (5.1%) reported that, during the 30 days prior to the survey, they had carried a weapon (such as a gun, knife, or club) while on school property. Overall, the prevalence for having carried such a weapon while on school property was higher among male students than female students (7.2% and 2.9%, respectively).

More than 1 in 20 students in surveyed SVVSD high schools (6.4%) reported that, in the 12 months before the survey, they had been threatened or injured with a weapon (such as a gun, knife, or club) while on school property. Overall, the prevalence of having been threatened or injured with such a weapon while on school property was higher among male students (8.5%) than female students (4.2%).

More than 1 in 10 students in surveyed SVVSD high schools (10.7%) reported they had been in a physical fight while on school property during the 12 months prior to the survey. Overall, the prevalence of having been in a physical fight on school property was higher among Hispanic students than white non-Hispanic students (15.5% and 8.9% respectively); it was also higher among male students than female students (14.3% and 6.9%, respectively).

Dating Violence and Electronic Bullying

	<i>Hurt by boyfriend/girlfriend*</i>			<i>Has been electronically bullied**</i>		
Grade	Total	Female	Male	Total	Female	Male
9th	5.2% (4.1-6.6)	7.4% (5.8-9.5)	3.0% (1.8-5.0)	17.1% (15.2-19.1)	28.0% (24.7-31.6)	6.4% (4.9-8.3)
10th	4.4% (3.5-5.5)	5.0% (3.8-6.7)	3.8% (2.7-5.3)	13.8% (12.0-15.8)	21.2% (18.2-24.6)	6.5% (4.7-8.9)
11th	9.6% (7.9-11.4)	8.7% (6.5-11.5)	10.4% (8.2-13.0)	16.1% (14.0-18.3)	24.3% (20.8-28.2)	8.4% (6.5-10.9)
12th	5.2% (4.0-6.7)	2.8% (1.7-4.5)	7.5% (5.5-10.1)	13.7% (11.3-16.4)	13.8% (10.5-17.9)	13.5% (10.5-17.3)
Race/Ethnicity						
White non-Hispanic	5.5% (4.8-6.3)	---	---	16.3% (14.9-17.7)	---	---
Hispanic	6.8% (5.5-8.5)	---	---	14.0% (12.1-16.0)	---	---
Other non-Hispanic	9.0% (6.5-12.3)	---	---	11.3% (8.4-15.0)	---	---
Total	6.1% (5.5-6.8)	6.0% (5.2-7.0)	6.2% (5.3-7.3)	15.2% (14.1-16.3)	21.9% (20.2-23.7)	8.7% (7.6-10.0)

* Hit, slapped, or physically hurt on purpose during the 12 months preceding the survey

** During the 12 months preceding the survey. Including via e-mail, chat rooms, instant messaging, websites, text messaging, and sexting. NOTE: The last portion of this question regarding sexting is not included on the 2009 National Survey.

--- Not available

Among students in surveyed SVVSD high schools, 6.1% reported they had been hit, slapped, or physically hurt on purpose by their boyfriend or girlfriend in the 12 months prior to the survey. Overall, the prevalence for having been hit, slapped, or physically hurt on purpose by their boyfriend or girlfriend was higher among other non-Hispanic students (9.0%) than white non-Hispanic students (5.5%).

Electronic bullying can occur via e-mail, chat rooms, instant messaging, websites, text messaging, or sexting (i.e., sexually explicit messages or photos sent by cell phone). Among students in surveyed SVVSD high schools, 15.2% reported having been electronically bullied at least once during the 12 months preceding the survey. Overall, the prevalence for having been electronically bullied was higher among female students than male students (21.9% and 8.7%, respectively).

Sexual Violence

Grade	<i>Ever forced to have sexual intercourse</i>			<i>Unwanted sexual contact*</i>		
	Total	Female	Male	Total	Female	Male
9th	4.4% (3.2-6.1)	7.0% (4.9-10.1)	1.9% (1.1-3.5)	14.2% (12.4-16.3)	23.5% (20.2-27.1)	5.3% (3.8-7.3)
10th	6.7% (5.3-8.5)	12.3% (9.7-15.5)	*** ***	13.0% (11.3-15.0)	23.2% (20.0-26.7)	3.1% (2.3-4.2)
11th	6.4% (5.1-8.0)	8.9% (6.8-11.5)	4.1% (2.7-6.4)	15.0% (13.0-17.2)	25.3% (21.8-29.0)	5.5% (3.8-8.1)
12th	8.5% (6.6-10.9)	8.7% (6.0-12.5)	8.4% (5.6-12.3)	16.7% (14.3-19.5)	23.1% (19.0-27.7)	10.6% (7.5-14.6)
Race/Ethnicity						
White non-Hispanic	5.2% (4.3-6.3)	---	---	15.2% (13.9-16.6)	---	---
Hispanic	8.7% (7.1-10.5)	---	---	14.1% (12.1-16.3)	---	---
Other non-Hispanic	10.0% (7.3-13.4)	---	---	13.8% (11.0-17.2)	---	---
Total	6.5% (5.7-7.4)	9.2% (7.9-10.7)	3.9% (3.0-5.1)	14.7% (13.7-15.8)	23.8% (22.0-25.7)	6.1% (5.1-7.4)

* Other than forced sexual intercourse

*** Numerator <3 and/or denominator <30

--- Not available

Among students in surveyed SVVSD high schools, 6.5% reported they had been forced to have sexual intercourse when they did not want to. Overall, the prevalence for being forced to have sex was higher among both other non-Hispanic and Hispanic students (10.0% and 8.7%, respectively) than white non-Hispanic students (5.2%); it was also higher among female students than male students (9.2% and 3.9%, respectively).

Nearly 1 in 7 students in surveyed SVVSD high schools (14.7%) reported they had been touched sexually when they did not want to be touched (excluding forced sexual intercourse). Overall, the prevalence for unwanted sexual contact was higher among female students (23.8%) than male students (6.1%). Additionally, at each of the four grade levels, female students had a higher prevalence of unwanted sexual contact than male students.



Sadness and Support

	<i>Felt sad or hopeless*</i>			<i>Has someone to talk to when feeling sad or hopeless</i>			<i>Usually gets help when feeling sad or hopeless</i>		
Grade	Total	Female	Male	Total	Female	Male	Total	Female	Male
9th	25.0% (22.7-27.5)	36.6% (32.7-40.6)	13.9% (11.4-16.9)	78.3% (75.8-80.6)	84.8% (81.5-87.7)	71.6% (67.9-74.9)	46.6% (43.8-49.4)	47.5% (43.6-51.5)	45.6% (41.8-49.5)
10th	21.2% (18.9-23.7)	27.8% (24.3-31.5)	14.8% (12.1-18.1)	81.2% (78.6-83.5)	90.5% (87.8-92.7)	71.3% (67.0-75.2)	47.5% (44.6-50.4)	54.0% (50.0-57.9)	40.6% (36.5-44.8)
11th	31.4% (28.8-34.1)	40.6% (36.5-44.8)	22.8% (19.7-26.3)	76.6% (73.8-79.2)	81.3% (77.2-84.7)	71.6% (67.4-75.4)	45.4% (42.4-48.4)	50.4% (46.1-54.6)	39.8% (35.6-44.1)
12th	27.4% (24.5-30.5)	36.9% (32.3-41.8)	18.1% (14.9-21.7)	85.3% (82.2-87.9)	87.6% (83.4-90.9)	82.4% (77.6-86.4)	50.4% (46.7-54.1)	52.3% (47.2-57.2)	48.0% (42.6-53.4)
Race/Ethnicity									
White non-Hispanic	25.9% (24.4-27.6)	---	---	83.0% (81.4-84.5)	---	---	50.3% (48.4-52.2)	---	---
Hispanic	27.4% (24.6-30.4)	---	---	78.8% (75.9-81.5)	---	---	46.8% (43.5-50.0)	---	---
Other non-Hispanic	25.3% (21.6-29.3)	---	---	66.7% (61.8-71.2)	---	---	32.4% (28.1-36.9)	---	---
Total	26.3% (25.0-27.6)	35.5% (33.5-37.6)	17.5% (15.9-19.1)	80.3% (79.0-81.6)	86.0% (84.3-87.6)	74.0% (71.9-75.9)	47.4% (45.9-49.0)	51.0% (48.8-53.2)	43.4% (41.2-45.6)

* Almost every day, 2 weeks or more, during the 12 months preceding the survey
 --- Not available

More than 1 in 4 students in surveyed SVVSD high schools (26.3%) reported that, in the 12 months before the survey, they had felt so sad or hopeless almost every day for 2 weeks or more that they had stopped doing usual activities. Overall, the prevalence for having felt sad or hopeless was higher among female students (35.5%) than male students (17.5%). Female students also had a higher prevalence for having felt sad or hopeless at each of the four grade levels than male students.

More than 4 in 5 students in surveyed SVVSD high schools (80.3%) reported they had someone they could talk with when they had felt sad, empty, hopeless, angry, or anxious. Overall, the prevalence for having someone they could talk to when they felt sad, empty, hopeless, angry, or anxious was higher among white non-Hispanic students (83.0%) and Hispanic students (78.8%), than other non-Hispanic students (66.7%). The prevalence was also higher among female students than male students (86.0% and 74.0%, respectively).

Nearly half of students in surveyed SVVSD high schools (47.4%) reported that when they felt sad, empty, hopeless, angry, or anxious, they got the help they needed always or most of the time. Overall, the prevalence for having gotten the help they needed was higher among both white non-Hispanic and Hispanic students (50.3% and 46.8%, respectively) than other non-Hispanic students (32.4%); it was also higher among female students (51.0%) than male students (43.4%).



Self-Injury and Suicide

	<i>Intentionally self-injured*</i>			<i>Seriously considered attempting suicide**</i>			<i>Planned suicide**</i>		
Grade	Total	Female	Male	Total	Female	Male	Total	Female	Male
9th	23.9% (21.7-26.3)	37.0% (33.3-40.8)	11.3% (9.3-13.7)	14.7% (12.8-16.8)	18.6% (15.5-22.0)	11.0% (8.8-13.7)	13.1% (11.3-15.2)	20.0% (16.8-23.7)	6.5% (5.1-8.2)
10th	18.8% (16.7-21.0)	24.6% (21.4-28.1)	13.1% (10.6-16.0)	13.3% (11.6-15.3)	16.3% (13.4-19.6)	10.5% (8.7-12.6)	9.5% (8.0-11.2)	12.8% (10.3-15.9)	6.3% (5.0-7.9)
11th	23.2% (21.0-25.7)	27.2% (23.7-30.9)	19.6% (16.8-22.7)	14.7% (12.7-16.9)	15.8% (12.8-19.4)	13.6% (11.1-16.5)	12.6% (10.9-14.7)	12.0% (9.5-15.1)	13.3% (10.9-16.1)
12th	15.3% (13.0-17.9)	17.9% (14.3-22.3)	12.7% (10.1-15.8)	11.5% (9.6-13.7)	13.2% (10.5-16.4)	9.9% (7.4-13.2)	9.2% (7.3-11.6)	9.6% (6.7-13.5)	8.9% (6.5-12.0)
Race/Ethnicity									
White non-Hispanic	20.0% (18.6-21.4)	---	---	13.9% (12.7-15.1)	---	---	11.0% (9.9-12.2)	---	---
Hispanic	20.7% (18.4-23.2)	---	---	11.5% (9.6-13.8)	---	---	10.5% (8.6-12.8)	---	---
Other non-Hispanic	21.6% (18.2-25.4)	---	---	17.0% (14.0-20.4)	---	---	13.1% (10.6-16.0)	---	---
Total	20.3% (19.2-21.5)	26.7% (24.9-28.6)	14.2% (12.9-15.6)	13.6% (12.6-14.6)	16.0% (14.5-17.6)	11.3% (10.1-12.6)	11.2% (10.2-12.2)	13.6% (12.2-15.3)	8.8% (7.7-9.9)

* Self-injured (such as cutting or burning) without wanting to die during the 12 months preceding the survey

** During the 12 months preceding the survey

--- Not available

More than 1 in 5 students in surveyed SVVSD high schools (20.3%) reported they had intentionally self-injured, such as cutting or burning, without wanting to die during the 12 months preceding the survey. Overall, the prevalence for intentional self-injury was higher among female students than male students (26.7% and 14.2%, respectively).

More than 1 in 10 students in surveyed SVVSD high schools (13.6%) reported they had seriously considered suicide during the 12 months prior to the survey. Overall, the prevalence for having seriously considered suicide was higher among other non-Hispanic students (17.0%) than Hispanic students (11.5%); it was also higher among female students (16.0%) than male students (11.3%).

More than 1 in 10 students in surveyed SVVSD high schools (11.2%) reported they had, during the 12 months preceding the survey, made a plan about how they would attempt suicide. Overall, the prevalence was higher among female students than male students (13.6% and 8.8%, respectively).



Suicide

Grade	Attempted suicide*			Suicide attempt required medical treatment*		
	Total	Female	Male	Total	Female	Male
9th	8.9% (7.1-11.0)	13.7% (10.7-17.2)	3.9% (2.2-6.9)	2.0% (1.1-3.7)	3.3% (1.6-6.6)	***
10th	7.4% (6.1-9.0)	9.0% (7.0-11.6)	5.7% (4.3-7.7)	2.3% (1.4-3.6)	3.7% (2.2-6.1)	***
11th	6.6% (5.2-8.4)	5.0% (3.2-7.7)	8.1% (6.1-10.8)	1.9% (1.2-3.0)	2.6% (1.4-4.8)	***
12th	2.7% (1.8-4.0)	4.2% (2.7-6.4)	***	***	***	***
Race/Ethnicity						
White non-Hispanic	5.4% (4.6-6.4)	---	---	1.7% (1.2-2.4)	---	---
Hispanic	9.9% (7.9-12.3)	---	---	2.0% (1.2-3.5)	---	---
Other non-Hispanic	4.9% (3.3-7.1)	---	---	***	---	---
Total	6.4% (5.7-7.2)	7.9% (6.8-9.3)	4.8% (3.9-5.9)	1.8% (1.4-2.4)	2.6% (1.9-3.6)	1.0% (0.6-1.5)

* During the 12 months preceding the survey
 *** Numerator <3 and/or denominator <30
 --- Not available

More than 1 in 20 students in surveyed SVVSD high schools (6.4%) reported they had attempted suicide at least once during the 12 months before the survey. Overall, the prevalence for attempted suicide was higher among Hispanic students (9.9%) than both other non-Hispanic and white non-Hispanic students (4.9% and 5.4%, respectively); it was also higher among female students (7.9%) than male students (4.8%).

Among students in surveyed SVVSD high schools, 1.8% reported a suicide attempt in the 12 months prior to the survey that resulted in an injury, poisoning, or overdose that required treatment by a doctor or nurse. Overall, the prevalence for having attempted suicide that required medical treatment was higher among female students than male students (2.6% and 1.0%, respectively).





Tobacco Use - Cigarettes

Grade	Ever smoked cigarettes*			Current cigarette user**			Current frequent cigarette use ⁺		
	Total	Female	Male	Total	Female	Male	Total	Female	Male
9th	31.4% (29.0-33.8)	27.5% (24.3-31.1)	35.1% (31.8-38.7)	13.9% (12.2-15.8)	15.7% (13.2-18.6)	12.2% (9.9-15.0)	***	***	***
10th	34.3% (31.7-37.0)	34.3% (30.6-38.2)	34.2% (30.4-38.2)	12.0% (10.3-14.0)	14.2% (11.6-17.3)	9.9% (7.8-12.4)	***	***	***
11th	45.6% (42.8-48.4)	48.3% (44.0-52.5)	43.1% (39.3-47.0)	17.9% (15.8-20.2)	17.2% (14.3-20.6)	18.5% (15.5-21.8)	7.3% (4.6-11.5)	***	***
12th	54.7% (51.5-58.0)	57.0% (52.1-61.7)	52.5% (47.8-57.2)	23.7% (20.9-26.7)	25.9% (21.5-30.9)	21.5% (17.8-25.7)	11.3% (6.4-19.0)	***	14.7% (9.1-22.8)
Race/Ethnicity									
White non-Hispanic	39.0% (37.3-40.7)	---	---	16.3% (15.0-17.7)	---	---	9.3% (6.2-13.8)	---	---
Hispanic	51.9% (48.9-54.9)	---	---	19.3% (17.0-21.9)	---	---	***	---	---
Other non-Hispanic	32.1% (27.7-36.8)	---	---	14.7% (11.5-18.7)	---	---	***	---	---
Total	41.5% (40.1-42.9)	41.7% (39.7-43.8)	41.3% (39.3-43.3)	16.9% (15.8-18.0)	18.3% (16.6-20.1)	15.6% (14.1-17.2)	6.9% (4.8-9.8)	5.5% (2.8-10.7)	8.5% (5.9-12.0)

* Even 1 or 2 puffs
 ** Smoked cigarettes on 1 or more of the 30 days preceding the survey
 + Among the 16.9% of students who reported current cigarette use
 *** Numerator <3 and/or denominator <30
 --- Not available

More than 2 in 5 students in surveyed SVVSD high schools (41.5%) reported they had ever tried smoking cigarettes, even if it was only 1 or 2 puffs. Overall, the prevalence for having tried smoking cigarettes was higher among Hispanic students (51.9%) than white non-Hispanic students (39.0%), and both were higher than other non-Hispanic students (32.1%).

Among students in surveyed SVVSD high schools, 16.9% reported current cigarette use (i.e., smoked cigarettes on at least 1 of the 30 days prior to the survey).

Among the 16.9% of students in surveyed SVVSD high schools who reported current cigarette use, 6.9% reported they smoked 10 or more cigarettes per day on the days they smoke.



Cigarettes - Purchase and Usage

	<i>Bought cigarettes in a store or gas station*</i>			<i>Ever smoked cigarettes daily**</i>			<i>Tried to quit smoking cigarettes⁺</i>		
Grade	Total	Female	Male	Total	Female	Male	Total	Female	Male
9th	*** ***	*** ***	*** ***	6.7% (5.4-8.2)	8.0% (6.2-10.4)	5.4% (3.8-7.5)	49.3% (42.6-56.1)	*** ***	*** ***
10th	*** ***	*** ***	*** ***	7.9% (6.5-9.5)	9.6% (7.4-12.3)	6.3% (4.7-8.3)	48.4% (40.2-56.5)	*** ***	*** ***
11th	7.5% (4.6-12.1)	*** ***	*** ***	11.6% (9.9-13.5)	9.6% (7.6-12.2)	13.3% (10.8-16.3)	49.5% (42.5-56.6)	*** ***	*** ***
12th	*** ***	*** ***	*** ***	12.5% (10.3-15.2)	12.4% (8.9-16.9)	12.6% (10.1-15.8)	43.2% (35.4-51.4)	53.4% (43.2-63.4)	*** ***
Race/Ethnicity									
White non-Hispanic	5.0% (3.2-7.6)	---	---	10.7% (9.5-11.9)	---	---	46.0% (41.0-51.1)	---	---
Hispanic	*** ***	---	---	8.2% (6.7-10.1)	---	---	46.9% (39.7-54.2)	---	---
Other non-Hispanic	*** ***	---	---	5.9% (4.0-8.5)	---	---	*** ***	---	---
Total	3.7% (2.5-5.6)	*** ***	7.6% (5.1-11.4)	9.7% (8.8-10.6)	9.9% (8.6-11.4)	9.4% (8.3-10.7)	47.1% (43.1-51.0)	50.6% (45.4-55.8)	43.0% (37.7-48.5)

* Among students who were less than 18 years old and report current cigarette use

** At least 1 cigarette every day for 30 days

+ In 12 months preceding the survey

*** Numerator <3 and/or denominator <30

--- Not available

Among students in surveyed SVVSD high schools who were younger than 18 years old and reported being a current cigarette user, 3.7% reported they had bought cigarettes in a convenience store, supermarket, discount store, or gas station during the 30 days prior to the survey.

Among students in surveyed SVVSD high schools, nearly 1 in 10 (9.7%) reported they had smoked at least 1 cigarette per day for 30 days (daily smoking). Overall, the prevalence of daily smoking was higher among white non-Hispanic students (10.7%) than other non-Hispanic students (5.9%).

Among the 47.1% of students in surveyed SVVD high schools who reported being a current cigarette user reported that, in the 12 months prior to the survey, they had tried to quit smoking.



Tobacco Use - Smokeless Tobacco, Cigars, and Overall Tobacco Use

Grade	Current smokeless tobacco use*			Current cigar use**			Current tobacco use*		
	Total	Female	Male	Total	Female	Male	Total	Female	Male
9th	6.3% (5.1-7.8)	2.0% (1.1-3.4)	10.6% (8.4-13.1)	10.7% (9.2-12.4)	11.6% (9.4-14.1)	9.8% (7.8-12.4)	17.9% (15.9-20.1)	16.1% (13.5-19.0)	19.6% (16.7-22.9)
10th	7.8% (6.5-9.5)	2.6% (1.5-4.6)	12.9% (10.6-15.6)	12.5% (10.7-14.5)	9.9% (7.9-12.4)	14.9% (12.2-18.1)	20.2% (17.9-22.4)	17.5% (14.6-20.7)	22.6% (19.4-26.1)
11th	9.3% (7.9-10.9)	4.3% (2.9-6.3)	14.0% (11.7-16.6)	18.0% (15.9-20.3)	12.3% (9.8-15.3)	23.2% (20.0-26.7)	27.2% (24.8-29.8)	20.9% (17.7-24.4)	33.1% (29.5-36.8)
12th	10.1% (8.4-12.2)	***	19.9% (16.6-23.7)	18.9% (16.2-21.9)	17.5% (13.6-22.3)	20.2% (16.9-24.0)	34.0% (30.7-37.3)	33.0% (28.2-38.2)	34.9% (30.5-39.6)
Race/Ethnicity									
White non-Hispanic	9.4% (8.5-10.5)	---	---	15.3% (13.9-16.7)	---	---	25.8% (24.2-27.4)	---	---
Hispanic	6.7% (5.3-8.5)	---	---	16.0% (14.0-18.2)	---	---	23.8% (21.3-26.5)	---	---
Other non-Hispanic	5.5% (3.9-7.8)	---	---	11.1% (8.5-14.4)	---	---	20.8% (17.1-25.0)	---	---
Total	8.4% (7.6-9.2)	2.2% (1.7-2.9)	14.3% (13.0-15.8)	15.0% (14.0-16.1)	12.8% (11.4-14.4)	17.1% (15.6-18.7)	24.8% (23.6-26.1)	21.8% (20.0-23.8)	27.6% (25.8-29.5)

* Used chewing tobacco or snuff on 1 or more of the 30 days preceding the survey

** Smoked cigars on 1 or more of the 30 days preceding the survey

+ Smoked cigarettes or cigars or used chewing tobacco or snuff on 1 or more of the 30 days preceding the survey

*** Numerator <3 and/or denominator <30

--- Not available

The prevalence of reported use of chewing tobacco, snuff, or dip during the 30 days prior to the survey among students in surveyed SVVSD high schools was 8.4%. Overall, the prevalence for having used smokeless tobacco was higher among white non-Hispanic students than other non-Hispanic students (9.4% and 5.5% respectively); it was also higher among male students than female students (14.3% and 2.2%, respectively).

Nearly 1 in 7 students in surveyed SVVSD high schools (15.0%) reported smoking cigars, cigarillos, or little cigars at least once during the 30 days prior to the survey. Overall, the prevalence for smoking cigars, cigarillos, or little cigars was higher in male students than female students (17.1% and 12.8%, respectively).

Nearly 1 in 4 students in surveyed SVVSD high schools (24.8%) reported they were current tobacco users (used at least once in the 30 days preceding the survey). Overall, the prevalence for being a current tobacco user was higher among male students (27.6%) than female students (21.8%).



Perceptions and Beliefs - Cigarettes

	<i>Smoking a pack/per day poses moderate to great risk</i>			<i>Wrong for someone your age to smoke cigarettes</i>			<i>Parents feel it is wrong for you to smoke cigarettes</i>		
Grade	Total	Female	Male	Total	Female	Male	Total	Female	Male
9th	88.1% (86.2-89.8)	87.6% (84.8-89.9)	88.7% (85.8-91.0)	73.8% (71.4-76.0)	75.5% (72.0-78.7)	72.0% (68.8-75.1)	95.9% (94.9-96.7)	97.1% (95.8-98.0)	94.7% (93.1-96.0)
10th	88.8% (86.9-90.5)	89.8% (87.2-92.0)	87.8% (84.9-90.2)	77.4% (75.0-79.7)	82.4% (79.1-85.2)	72.5% (68.9-75.9)	92.2% (90.5-93.7)	92.9% (90.4-94.8)	91.6% (88.9-93.6)
11th	89.8% (87.9-91.4)	91.6% (89.0-93.6)	88.1% (85.2-90.5)	64.5% (61.7-67.2)	63.1% (59.1-67.0)	65.8% (62.0-69.3)	90.5% (88.9-92.0)	93.6% (91.8-95.0)	87.7% (85.0-90.0)
12th	91.1% (89.3-92.6)	95.9% (94.1-97.2)	86.4% (83.1-89.1)	59.2% (55.8-62.6)	58.4% (53.3-63.4)	60.0% (55.4-64.4)	90.7% (88.4-92.7)	94.3% (90.7-96.6)	87.3% (83.9-90.1)
Race/Ethnicity									
White non-Hispanic	92.4% (91.4-93.2)	---	---	67.8% (66.1-69.5)	---	---	92.4% (91.3-93.3)	---	---
Hispanic	83.8% (81.4-85.9)	---	---	71.9% (69.1-74.6)	---	---	92.4% (90.7-93.8)	---	---
Other non-Hispanic	86.7% (83.3-89.5)	---	---	66.1% (61.6-70.3)	---	---	93.1% (90.7-94.9)	---	---
Total	89.4% (88.5-90.3)	91.2% (90.0-92.3)	87.7% (86.3-89.0)	68.7% (67.3-70.0)	69.9% (67.9-71.9)	67.5% (65.6-69.4)	92.4% (91.5-93.1)	94.5% (93.4-95.4)	90.3% (89.0-91.4)

--- Not available

Nearly 9 in 10 students in surveyed SVVSD high schools (89.4%) reported they believed smoking 1 or more packs of cigarettes per day posed moderate to great risk of harm. Overall, prevalence of that belief was higher among white non-Hispanic students (92.4%) than both other non-Hispanic and Hispanic students (86.7% and 83.8%, respectively); it was also higher in female students (91.2%) than male students (87.7%).

Among students in surveyed SVVSD high schools, 68.7% reported they believed that it was wrong or very wrong for someone their own age to smoke cigarettes.

More than 9 in 10 students in surveyed SVVSD high schools (92.4%) reported they believed their parents/guardians felt it was wrong or very wrong for them to smoke cigarettes. Overall, prevalence of this belief was higher among female students (94.5%) than male students (90.3%).



Alcohol Use

	<i>Ever drank alcohol*</i>			<i>Current alcohol use**</i>			<i>Binge drinking⁺</i>		
Grade	Total	Female	Male	Total	Female	Male	Total	Female	Male
9th	55.1% (52.3-57.8)	56.9% (52.7-61.0)	53.4% (49.5-57.2)	30.9% (28.6-33.3)	35.4% (31.9-39.1)	26.3% (23.2-29.6)	14.7% (13.0-16.5)	20.3% (17.6-23.4)	9.0% (7.1-11.5)
10th	62.5% (59.8-65.1)	66.6% (62.7-70.2)	58.7% (54.8-62.5)	37.3% (34.7-40.0)	40.1% (36.2-44.1)	34.6% (30.9-38.6)	24.7% (22.3-27.1)	23.4% (20.1-27.1)	25.8% (22.4-29.6)
11th	77.4% (74.9-79.7)	83.3% (79.8-86.2)	71.6% (67.8-75.0)	42.8% (40.0-45.6)	46.6% (42.4-50.9)	39.3% (35.6-43.2)	25.8% (23.4-28.3)	26.0% (22.6-29.8)	25.5% (22.3-29.1)
12th	82.5% (79.7-84.9)	86.4% (82.9-89.3)	78.6% (74.1-82.5)	51.8% (48.4-55.2)	55.8% (50.8-60.7)	47.9% (43.0-52.8)	37.3% (34.0-40.8)	38.8% (34.0-43.9)	35.8% (31.3-40.5)
Race/Ethnicity									
White non-Hispanic	68.1% (66.4-69.7)	---	---	40.3% (38.5-42.1)	---	---	25.9% (24.3-27.5)	---	---
Hispanic	77.6% (74.8-80.2)	---	---	46.3% (43.2-49.3)	---	---	28.4% (25.6-31.3)	---	---
Other non-Hispanic	62.7% (58.1-67.1)	---	---	31.3% (27.1-35.8)	---	---	16.4% (13.3-19.9)	---	---
Total	69.7% (68.4-71.0)	73.8% (72.0-75.6)	65.7% (63.8-67.7)	40.7% (39.3-42.1)	44.5% (42.3-46.6)	37.1% (35.1-39.1)	25.5% (24.2-26.8)	27.1% (25.2-29.1)	24.0% (22.2-25.8)

* More than a few sips

** Drank at least 1 drink of alcohol on 1 or more of the 30 days preceding the survey

+ Drank 5 or more drinks of alcohol in a row (within a couple of hours) on 1 or more of the 30 days preceding the survey

--- Not available

Nearly 7 in 10 students in surveyed SVVSD high schools (69.7%) reported ever having had more than a few sips of alcohol (e.g., beer, wine, wine coolers, and liquor) in their life. Overall, the prevalence of alcohol use was higher among Hispanic students (77.6%) than both white non-Hispanic students and other non-Hispanic students (68.1% and 62.7%, respectively); it was also higher among female students (73.8%) than male students (65.7%).

More than 2 in 5 students in surveyed SVVSD high schools (40.7%) reported they were current alcohol users (i.e., had drunk alcohol on at least 1 of the 30 days prior to the survey). Overall, prevalence of current alcohol use was higher among Hispanic students (46.3%) than white non-Hispanic students (40.3%), and both were higher than other non-Hispanic students (31.3%). The prevalence was also higher among female students than male students (44.5% and 37.1%, respectively).

Binge drinking (defined as having 5 or more drinks of alcohol in a row within a few hours) during the 30 days prior to the survey was reported by 1 in 4 students in surveyed SVVSD high schools (25.5%). Overall, the prevalence of binge drinking was higher among both Hispanic and white non-Hispanic students (28.4% and 25.9%, respectively) than other non-Hispanic students (16.4%).



Perceptions and Beliefs - Alcohol

	<i>1 or 2 drinks/day poses moderate to great risk</i>			<i>Wrong for someone your age to drink</i>			<i>Parents feel it is wrong for you to drink</i>		
Grade	Total	Female	Male	Total	Female	Male	Total	Female	Male
9th	62.4% (59.7-65.0)	69.8% (66.1-73.3)	55.0% (51.2-58.7)	72.6% (70.1-75.0)	74.4% (70.8-77.6)	70.9% (67.3-74.2)	88.6% (86.8-90.1)	90.7% (88.3-92.6)	86.5% (83.7-88.8)
10th	62.1% (59.3-64.7)	70.5% (66.8-73.9)	53.9% (49.9-57.7)	69.3% (66.6-71.9)	76.2% (72.7-79.4)	62.6% (58.7-66.4)	89.7% (87.8-91.3)	94.2% (92.0-95.8)	85.3% (82.2-88.0)
11th	60.5% (57.6-63.3)	66.5% (62.3-70.4)	54.8% (50.8-58.7)	60.9% (58.1-63.7)	60.8% (56.6-65.0)	60.9% (57.1-64.6)	82.1% (79.9-84.2)	83.4% (80.0-86.3)	80.9% (77.9-83.7)
12th	56.1% (52.7-59.5)	60.4% (55.5-65.2)	51.9% (47.2-56.7)	58.2% (54.8-61.5)	59.2% (54.2-64.1)	57.2% (52.5-61.8)	81.0% (78.1-83.6)	82.6% (78.1-86.3)	79.4% (75.4-82.9)
Race/Ethnicity									
White non-Hispanic	58.1% (56.3-59.9)	---	---	64.6% (62.9-66.3)	---	---	83.8% (82.4-85.1)	---	---
Hispanic	63.2% (60.1-66.2)	---	---	65.5% (62.4-68.4)	---	---	88.0% (85.9-89.8)	---	---
Other non-Hispanic	66.0% (61.5-70.2)	---	---	67.6% (63.0-72.0)	---	---	89.2% (85.7-91.9)	---	---
Total	60.3% (58.8-61.7)	66.8% (64.8-68.8)	53.9% (51.9-56.0)	65.3% (63.8-66.7)	67.7% (65.6-69.7)	62.9% (60.9-64.9)	85.3% (84.2-86.3)	87.7% (86.1-89.1)	83.0% (81.4-84.5)

--- Not available

More than 3 in 5 students in surveyed SVVSD high schools (60.3%) reported the belief that consumption of 1 or 2 drinks of alcohol nearly every day posed a moderate or great risk for harm to themselves. Overall, the prevalence of that belief was higher among both other non-Hispanic and Hispanic students (66.0% and 63.2%, respectively) than white non-Hispanic students (58.1%); it was also higher among female students (66.8%) than male students (53.9%).

Among students in surveyed SVVSD high schools, 65.3% reported it was wrong or very wrong for someone their age to drink beer, wine, or hard liquor regularly. Overall, the prevalence of this belief was higher among female students than male students (67.7% and 62.9%, respectively).

More than 4 in 5 students in surveyed SVVSD high schools (85.3%) reported they believed their parents/guardians felt it was wrong or very wrong for them to drink beer, wine, or hard liquor regularly. Overall, the prevalence of this belief was higher among both other non-Hispanic and Hispanic students (89.2% and 88.0%, respectively) than white non-Hispanic students (83.8%); it was also higher among female students (87.7%) than male students (83.0%).



Marijuana Use

Grade	Ever used marijuana			Current marijuana use*		
	Total	Female	Male	Total	Female	Male
9th	25.1% (22.8-27.5)	21.9% (18.7-25.4)	28.2% (25.0-31.7)	15.3% (13.5-17.2)	13.0% (10.7-15.6)	17.5% (14.9-20.5)
10th	33.8% (31.3-36.4)	27.4% (24.0-31.0)	40.0% (36.3-43.9)	19.0% (16.9-21.3)	16.9% (14.1-20.2)	21.0% (18.0-24.3)
11th	42.9% (40.1-45.9)	48.8% (44.5-53.1)	37.5% (33.7-41.5)	22.8% (20.4-25.3)	22.1% (18.8-25.8)	23.4% (20.1-27.0)
12th	48.4% (45.1-51.7)	49.9% (45.0-54.8)	46.9% (42.3-51.7)	23.8% (21.0-26.9)	25.3% (20.9-30.2)	22.4% (18.6-26.7)
Race/Ethnicity						
White non-Hispanic	35.9% (34.2-37.6)	---	---	20.7% (19.3-22.3)	---	---
Hispanic	45.9% (42.8-49.1)	---	---	21.6% (19.2-24.3)	---	---
Other non-Hispanic	28.3% (24.1-33.0)	---	---	14.3% (11.1-18.2)	---	---
Total	37.6% (36.2-39.0)	37.0% (34.9-39.1)	38.2% (36.2-40.2)	20.2% (19.1-21.5)	19.3% (17.6-21.2)	21.1% (19.5-22.9)

* Used marijuana 1 or more times during the 30 days preceding the survey

--- Not available

Among students in surveyed SVVSD high schools, 37.6% reported having ever used marijuana in their life. Overall, the prevalence of marijuana use was higher among Hispanic students (45.9%) than white non-Hispanic students (35.9%), and both were higher than other non-Hispanic students (28.3%).

More than 1 in 5 students in surveyed SVVSD high schools (20.2%) reported they were current marijuana users (i.e., had used marijuana on at least 1 of the 30 days preceding the survey). Overall, the prevalence of current marijuana use was higher among both Hispanic and white non-Hispanic students (21.6% and 20.7%, respectively) than other non-Hispanic students (14.3%).



Perceptions and Beliefs - Marijuana

	<i>Smoking marijuana regularly poses moderate to great risk</i>			<i>Wrong for someone your age to smoke marijuana</i>			<i>Parents feel it is wrong for you to smoke marijuana</i>		
Grade	Total	Female	Male	Total	Female	Male	Total	Female	Male
9th	68.7% (66.3-71.1)	71.3% (67.7-74.7)	66.2% (62.8-69.5)	70.0% (67.4-72.5)	73.0% (69.2-76.5)	67.0% (63.4-70.4)	94.1% (92.8-95.2)	94.2% (92.1-95.7)	94.0% (92.3-95.3)
10th	67.4% (64.8-69.9)	73.0% (69.5-76.3)	61.9% (58.1-65.5)	69.4% (66.8-71.9)	74.1% (70.7-77.3)	64.9% (61.1-68.6)	91.2% (89.4-92.7)	89.4% (86.7-91.6)	92.9% (90.4-94.8)
11th	54.8% (52.0-57.6)	56.3% (52.1-60.5)	53.5% (49.6-57.3)	53.9% (51.0-56.7)	49.5% (45.3-53.6)	57.9% (54.0-61.7)	87.1% (85.1-88.9)	87.1% (84.3-89.5)	87.0% (84.1-89.5)
12th	55.9% (52.4-59.4)	57.6% (52.5-62.5)	54.2% (49.4-59.0)	51.0% (47.6-54.4)	46.8% (41.9-51.8)	55.1% (50.3-59.7)	90.0% (87.6-92.0)	89.3% (85.7-92.2)	90.7% (87.2-93.2)
Race/Ethnicity									
White non-Hispanic	62.0% (60.3-63.8)	---	---	58.7% (57.0-60.5)	---	---	90.4% (89.3-91.5)	---	---
Hispanic	60.6% (57.6-63.5)	---	---	65.6% (62.5-68.5)	---	---	90.4% (88.5-92.0)	---	---
Other non-Hispanic	61.6% (57.0-66.1)	---	---	63.2% (58.7-67.6)	---	---	92.2% (89.0-94.6)	---	---
Total	61.6% (60.2-63.1)	64.5% (62.4-66.6)	58.9% (56.9-60.8)	61.0% (59.6-62.4)	60.9% (58.8-62.9)	61.2% (59.2-63.2)	90.6% (89.7-91.4)	90.0% (88.7-91.3)	91.1% (89.8-92.2)

--- Not available

More than 3 in 5 students in surveyed SVVSD high schools (61.6%) reported they believed that regularly smoking marijuana posed a moderate or great risk. Overall, the prevalence of this belief was higher among female students than male students (64.5% and 58.9%, respectively).

More than 3 in 5 students in surveyed SVVSD high schools (61.0%) reported they believed it was wrong or very wrong for someone their age to smoke marijuana. Overall, the prevalence of this belief was higher among Hispanic students (65.6%) than white non-Hispanic students (58.7%).

More than 9 in 10 students in surveyed SVVSD high schools (90.6%) reported they believed their parents/guardians thought it was wrong or very wrong for them to smoke marijuana.



Early Initiation of Tobacco, Alcohol, and Marijuana Use

	<i>Smoked a whole cigarette before age 13</i>			<i>Drank alcohol before age 13*</i>			<i>Smoked marijuana before age 13</i>		
Grade	Total	Female	Male	Total	Female	Male	Total	Female	Male
9th	11.1% (9.6-12.8)	9.3% (7.4-11.8)	12.8% (10.6-15.4)	29.3% (26.8-31.9)	28.9% (25.4-32.8)	29.6% (26.1-33.4)	8.2% (6.9-9.7)	6.5% (4.8-8.6)	9.9% (8.0-12.2)
10th	8.8% (7.4-10.6)	6.9% (5.1-9.1)	10.7% (8.5-13.5)	22.8% (20.3-25.4)	22.5% (19.1-26.4)	23.0% (19.7-26.6)	7.3% (6.2-8.7)	6.4% (5.0-8.3)	8.2% (6.5-10.4)
11th	7.5% (6.1-9.3)	7.2% (5.2-9.8)	7.9% (5.9-10.5)	17.6% (15.4-20.1)	15.4% (12.5-18.9)	19.8% (16.7-23.4)	6.2% (4.8-7.8)	4.3% (2.8-6.4)	7.9% (5.8-10.6)
12th	9.1% (7.2-11.4)	6.3% (3.9-9.9)	11.7% (9.1-15.0)	16.6% (14.0-19.6)	11.9% (8.6-16.4)	21.2% (17.3-25.8)	6.1% (4.6-8.0)	3.7% (1.9-7.2)	8.4% (6.4-11.0)
Race/Ethnicity									
White non-Hispanic	7.4% (6.5-8.5)	---	---	19.0% (17.5-20.6)	---	---	5.7% (4.9-6.6)	---	---
Hispanic	15.1% (13.0-17.4)	---	---	27.9% (25.1-30.8)	---	---	10.9% (9.2-12.8)	---	---
Other non-Hispanic	4.4% (3.0-6.3)	---	---	20.0% (16.3-24.2)	---	---	3.7% (2.1-6.6)	---	---
Total	9.1% (8.3-10.0)	7.4% (6.3-8.7)	10.8% (9.6-12.1)	21.4% (20.2-22.8)	19.4% (17.7-21.3)	23.4% (21.5-25.3)	6.9% (6.2-7.7)	5.2% (4.3-6.3)	8.6% (7.6-9.8)

* More than a few sips

--- Not available

Among students in surveyed SVVSD high schools, 9.1% reported they had smoked a whole cigarette before the age of 13. Overall, the prevalence of having smoked a whole cigarette before age 13 was higher among Hispanic students (15.1%) than white non-Hispanic students (7.4%), and both were higher than other non-Hispanic students (4.4%). The prevalence was higher among male students (10.8%) than female students (7.4%).

More than 1 in 5 students in surveyed SVVSD high schools (21.4%) reported they had had their first drink of alcohol (other than a few sips) before the age of 13. Overall, the prevalence of having their first drink of alcohol before age 13 was higher among Hispanic students (27.9%) than both other non-Hispanic and white non-Hispanic students (20.0% and 19.0%, respectively); it was also higher among male students (23.4%) than female students (19.4%).

Among students in surveyed SVVSD high schools, 6.9% reported they had tried marijuana before the age of 13. Overall, the prevalence of having tried marijuana before the age of 13 was higher among Hispanic students (10.9%) than both white non-Hispanic and other non-Hispanic students (5.7% and 3.7%, respectively); it was also higher among male students (8.6%) than female students (5.2%).



Drug Use - Cocaine, Inhalants, and Ecstasy

	<i>Ever used cocaine*</i>			<i>Ever used inhalants**</i>			<i>Ever used ecstasy</i>		
Grade	Total	Female	Male	Total	Female	Male	Total	Female	Male
9th	6.2% (5.0-7.6)	7.0% (5.4-9.1)	5.3% (3.8-7.4)	11.9% (10.3-13.8)	16.1% (13.6-19.0)	7.8% (5.9-10.2)	6.4% (5.3-7.8)	9.1% (7.2-11.6)	3.8% (2.7-5.3)
10th	5.8% (4.6-7.3)	5.3% (3.9-7.2)	6.3% (4.5-8.7)	11.6% (9.8-13.6)	12.3% (9.6-15.5)	10.9% (8.5-13.8)	9.7% (8.0-11.6)	11.3% (8.9-14.3)	8.1% (6.0-10.8)
11th	11.0% (9.4-13.0)	10.3% (8.0-13.0)	11.8% (9.3-14.7)	9.7% (8.0-11.7)	11.0% (8.4-14.3)	8.6% (6.5-11.1)	15.5% (13.5-17.7)	16.2% (13.5-19.4)	14.8% (12.1-17.9)
12th	8.5% (6.7-10.8)	6.0% (3.6-9.8)	11.0% (8.5-14.1)	8.1% (6.5-10.0)	7.6% (5.2-10.9)	8.5% (6.7-10.9)	14.2% (12.0-16.9)	14.9% (11.2-19.6)	13.6% (10.4-17.4)
Race/Ethnicity									
White non-Hispanic	6.1% (5.2-7.1)	---	---	9.6% (8.6-10.8)	---	---	10.9% (9.8-12.2)	---	---
Hispanic	13.0% (11.1-15.1)	---	---	12.0% (10.2-14.1)	---	---	12.7% (10.8-14.9)	---	---
Other non-Hispanic	6.8% (4.7-9.8)	---	---	9.9% (7.5-13.0)	---	---	11.2% (8.5-14.7)	---	---
Total	7.9% (7.1-8.8)	7.2% (6.1-8.4)	8.6% (7.5-9.9)	10.3% (9.5-11.3)	11.8% (10.4-13.2)	8.9% (7.9-10.1)	11.5% (10.5-12.5)	12.9% (11.4-14.5)	10.1% (8.9-11.5)

* Ever tried any form of cocaine (e.g., "powder," "crack," "freebase," "blow," or "yah")

** Ever sniffed glue or breathed the contents of aerosol spray cans or inhaled any paints to get high

--- Not available

Among students in surveyed SVVSD high schools, 7.9% reported they had ever used any form of cocaine (e.g., powder, crack, freebase, blow, or yah). Overall, the prevalence of ever having used any form of cocaine was higher among Hispanic students (13.0%) than both other non-Hispanic and white non-Hispanic students (6.8% and 6.1%, respectively).

More than 1 in 10 students in surveyed SVVSD high schools (10.3%) reported they had ever dusted, huffed, sniffed, inhaled, or breathed the contents of aerosol spray cans, glues, or any paints or sprays to get high. Overall, the prevalence of having used inhalants was higher among female students than male students (11.8% and 8.9%, respectively).

More than 1 in 10 students in surveyed SVVSD high schools (11.5%) reported they had ever used ecstasy (also called MDMA, X, candy, molly, E, rolling).



Drug Use - Heroin, Methamphetamine, and Needle Use

Grade	Ever used heroin*			Ever used methamphetamine*			Ever used a needle to inject drugs*		
	Total	Female	Male	Total	Female	Male	Total	Female	Male
9th	1.9% (1.2-2.8)	2.0% (1.2-3.3)	1.8% (1.0-3.2)	2.8% (2.0-3.9)	4.2% (2.8-6.3)	1.4% (0.9-2.1)	1.8% (1.2-2.7)	2.6% (1.6-4.4)	*** ***
10th	2.6% (1.7-3.8)	*** ***	3.5% (2.3-5.2)	2.6% (1.7-3.8)	2.4% (1.2-4.7)	2.7% (1.7-4.3)	1.8% (1.1-2.9)	*** ***	2.1% (1.2-3.4)
11th	2.0% (1.4-2.8)	*** ***	3.2% (2.1-4.6)	2.7% (1.9-3.7)	*** ***	4.5% (3.1-6.5)	1.9% (1.4-2.8)	*** ***	3.8% (2.6-5.3)
12th	2.2% (1.5-3.2)	*** ***	4.3% (2.9-6.3)	3.2% (2.0-5.0)	*** ***	4.2% (2.8-6.2)	1.2% (0.7-2.1)	*** ***	2.3% (1.3-4.1)
Race/Ethnicity									
White non-Hispanic	1.7% (1.3-2.3)	--- ---	--- ---	2.4% (1.8-3.2)	--- ---	--- ---	1.5% (1.2-2.0)	--- ---	--- ---
Hispanic	3.2% (2.3-4.3)	--- ---	--- ---	4.1% (3.1-5.5)	--- ---	--- ---	2.5% (1.7-3.6)	--- ---	--- ---
Other non-Hispanic	*** ***	--- ---	--- ---	*** ***	--- ---	--- ---	*** ***	--- ---	--- ---
Total	2.1% (1.8-2.6)	1.1% (0.7-1.6)	3.2% (2.6-3.9)	2.8% (2.3-3.4)	2.4% (1.6-3.4)	3.2% (2.6-4.0)	1.7% (1.4-2.1)	1.1% (0.7-1.7)	2.3% (1.8-2.9)

* One or more times in your life

*** Numerator <3 and/or denominator <30

--- Not available

Among students in surveyed SVVSD high schools, 2.1% reported they had ever used heroin (also called smack, junk, or China White). Overall, the prevalence of having used heroin was higher among male students than female students (3.2% and 1.1% respectively).

Among students in surveyed SVVSD high schools, 2.8% reported they had ever used methamphetamines (also called speed, crystal, crank, ice, meth, or tweaking).

Among students in surveyed SVVSD high schools, 1.7% reported they had ever used a needle to inject any illegal drug into their body. Overall, the prevalence of ever using a needle to inject any illegal drugs was higher among male students than female students (2.3% and 1.1%, respectively).

Drug Use - Illegal Steroids and Prescription Drugs

Grade	Ever used illegal steroids*			Ever used prescription drugs without a prescription**		
	Total	Female	Male	Total	Female	Male
9th	2.8% (2.1-3.7)	2.4% (1.6-3.6)	3.2% (2.2-4.6)	12.2% (10.6-13.9)	14.8% (12.3-17.7)	9.6% (7.7-11.8)
10th	3.8% (2.9-4.8)	4.4% (3.1-6.3)	3.1% (2.1-4.5)	15.3% (13.4-17.5)	18.1% (15.1-21.4)	12.7% (10.3-15.6)
11th	2.9% (2.2-3.9)	4.8% (3.4-6.8)	*** ***	23.3% (20.9-25.9)	22.7% (19.3-26.5)	23.9% (20.6-27.5)
12th	3.4% (2.5-4.4)	1.9% (1.2-3.0)	4.8% (3.4-6.7)	20.4% (17.8-23.3)	20.0% (16.0-24.6)	20.9% (17.2-25.1)
Race/Ethnicity						
White non-Hispanic	3.1% (2.6-3.7)	---	---	18.4% (17.0-19.8)	---	---
Hispanic	3.1% (2.3-4.1)	---	---	17.2% (15.0-19.7)	---	---
Other non-Hispanic	3.9% (2.5-6.0)	---	---	17.2% (13.9-21.0)	---	---
Total	3.2% (2.8-3.7)	3.4% (2.8-4.1)	3.0% (2.5-3.7)	17.9% (16.8-19.0)	18.9% (17.2-20.7)	16.9% (15.4-18.5)

* Ever taken steroid pills or shots without a doctor's prescription
 ** Ever used prescription drugs without a doctor's prescription (e.g., OxyContin, Percocet, Vicodin, Adderall, Ritalin, or Xanax)
 *** Numerator <3 and/or denominator <30
 --- Not available

Less than 1 in 25 students in surveyed SVVSD high schools (3.2%) reported they had ever taken steroid pills or shots without a doctor's prescription.

More than 1 in 5 students in surveyed SVVSD high schools (17.9%) reported they had ever taken a prescription drug (e.g., OxyContin, Percocet, Vicodin, Adderall, Ritalin, or Xanax) without a doctor's prescription.

Other Drug Use and Current Cocaine Use

	<i>Ever used other illegal drug*</i>			<i>Current cocaine use**</i>		
Grade	Total	Female	Male	Total	Female	Male
9th	7.4% (6.0-9.1)	10.0% (7.6-13.1)	4.8% (3.6-6.4)	4.1% (3.2-5.4)	5.2% (3.8-7.0)	3.2% (1.9-5.1)
10th	10.2% (8.6-12.1)	8.9% (6.8-11.6)	11.4% (9.0-14.4)	2.4% (1.8-3.2)	3.0% (2.0-4.4)	1.9% (1.2-2.9)
11th	12.6% (10.9-14.7)	8.7% (6.8-11.2)	16.3% (13.5-19.5)	5.4% (4.2-6.9)	3.5% (2.3-5.5)	7.1% (5.3-9.5)
12th	13.0% (10.8-15.5)	11.5% (8.3-15.6)	14.4% (11.1-18.5)	1.9% (1.3-2.9)	***	2.9% (1.8-4.7)
Race/Ethnicity						
White non-Hispanic	10.0% (8.9-11.2)	---	---	2.6% (2.1-3.2)	---	---
Hispanic	13.5% (11.5-15.8)	---	---	6.0% (4.8-7.5)	---	---
Other non-Hispanic	8.4% (6.1-11.3)	---	---	2.8% (1.5-5.2)	---	---
Total	10.8% (9.9-11.8)	9.8% (8.5-11.3)	11.8% (10.4-13.2)	3.5% (3.0-4.0)	3.1% (2.5-3.9)	3.8% (3.1-4.7)

* Ever used any other type of illegal drugs, such as LSD (acid), PCP, mushrooms, Ketamine (Special K), Rohypnol (Roofies), or GHB

** During the 30 days preceding the survey, including any form of cocaine

*** Numerator <3 and/or denominator <30

--- Not available

Other types of illegal drugs include LSD (acid), phencyclidine (PCP), mushrooms (shrooms), Ketamine (Special K), Rohypnol (Roofies), or GHB (liquid X or liquid ecstasy). More than 1 in 10 students in surveyed SVVSD high schools (10.8%) reported ever having used any of these drugs. Overall, the prevalence of ever using other drugs was higher among Hispanic students (13.5%) than both white non-Hispanic and other non-Hispanic students (10.0% and 8.4%, respectively).

Among students in surveyed SVVSD high schools, 3.5% reported they were current cocaine users (i.e., used some form of cocaine on at least 1 of the 30 days prior to the survey). Overall, the prevalence of current cocaine use was higher among Hispanic students (6.0%) than white non-Hispanic students (2.6%).

Drug and Alcohol Activity on School Property

	<i>Offered, sold, or given an illegal drug on school property*</i>			<i>Attended school under the influence of alcohol or other illegal drugs*</i>		
Grade	Total	Female	Male	Total	Female	Male
9th	21.4% (19.2-23.7)	21.2% (18.1-24.8)	21.5% (18.4-24.9)	13.8% (12.1-15.6)	14.8% (12.3-17.8)	12.7% (10.6-15.2)
10th	16.7% (14.7-19.0)	15.1% (12.3-18.4)	18.3% (15.4-21.6)	15.5% (13.5-17.8)	15.2% (12.4-18.6)	15.8% (13.0-19.0)
11th	23.1% (20.8-25.6)	18.7% (15.8-22.1)	27.2% (23.7-30.9)	20.4% (18.1-22.8)	20.7% (17.5-24.4)	20.0% (17.0-23.5)
12th	17.0% (15.0-19.3)	15.6% (12.8-18.9)	18.4% (15.4-21.9)	15.7% (13.5-18.2)	13.0% (10.2-16.6)	18.4% (15.1-22.2)
Race/Ethnicity						
White non-Hispanic	17.9% (16.6-19.3)	---	---	14.7% (13.5-16.0)	---	---
Hispanic	24.4% (21.8-27.2)	---	---	21.6% (19.1-24.4)	---	---
Other non-Hispanic	20.0% (16.6-23.8)	---	---	14.9% (12.0-18.4)	---	---
Total	19.6% (18.5-20.8)	17.7% (16.2-19.3)	21.4% (19.8-23.2)	16.4% (15.3-17.5)	15.9% (14.4-17.6)	16.8% (15.3-18.4)

* During the 12 months preceding the survey
 --- Not available

Nearly 1 in 5 students in surveyed SVVSD high schools (19.6%) reported they had been offered, sold, or given an illegal drug while on school property during the 12 months prior to the survey. Overall, the prevalence of this behavior was higher among Hispanic students (24.4%) than white non-Hispanic students (17.9%); it was also higher among male students (21.4%) than female students (17.7%).

More than 1 in 7 students in surveyed SVVSD high schools (16.4%) reported that, in the 12 months preceding the survey, they had attended school at least once while under the influence of alcohol or other illegal drugs. Overall, the prevalence of having attended school while under the influence was higher among Hispanic students (21.6%) than both other non-Hispanic and white non-Hispanic students (14.9% and 14.7%, respectively).

Substance Use on School Property - Cigarettes, Smokeless Tobacco, Alcohol

Grade	Smoked cigarettes on school property*			Used smokeless tobacco on school property**			Drank alcohol on school property*		
	Total	Female	Male	Total	Female	Male	Total	Female	Male
9th	4.0% (3.1-5.1)	4.4% (3.2-6.0)	3.6% (2.4-5.3)	3.9% (3.0-5.1)	***	6.6% (4.9-8.7)	2.6% (1.8-3.7)	3.2% (2.1-4.8)	1.9% (1.0-3.7)
10th	5.4% (4.3-6.7)	6.8% (5.1-8.9)	4.0% (2.7-5.9)	2.4% (1.7-3.3)	***	4.2% (2.9-5.9)	2.2% (1.5-3.5)	2.9% (1.8-4.8)	***
11th	7.0% (5.7-8.6)	4.7% (3.3-6.8)	9.1% (7.0-11.8)	5.3% (4.3-6.5)	***	10.2% (8.3-12.6)	4.4% (3.3-5.9)	***	7.8% (5.8-10.4)
12th	7.6% (5.8-9.8)	5.0% (2.8-8.9)	10.1% (7.7-13.1)	6.9% (5.6-8.7)	***	13.7% (11.0-16.9)	3.3% (2.4-4.5)	***	6.5% (4.7-8.9)
Race/Ethnicity									
White non-Hispanic	6.1% (5.3-7.1)	---	---	5.8% (5.0-6.6)	---	---	2.8% (2.3-3.5)	---	---
Hispanic	6.3% (4.9-8.0)	---	---	3.1% (2.2-4.3)	---	---	4.3% (3.1-5.8)	---	---
Other non-Hispanic	4.1% (2.6-6.3)	---	---	***	---	---	***	---	---
Total	6.0% (5.3-6.7)	5.2% (4.3-6.3)	6.7% (5.8-7.8)	4.6% (4.1-5.2)	0.4% (0.3-0.7)	8.7% (7.7-9.8)	3.1% (2.7-3.7)	1.7% (1.3-2.4)	4.5% (3.7-5.5)

* During the 30 days preceding the survey

** During the 30 days preceding the survey; including chewing tobacco, snuff, or dip

*** Numerator <3 and/or denominator <30

--- Not available

More than 1 in 20 students in surveyed SVVSD high schools (6.0%) reported they had smoked a cigarette while on school property during the 30 days prior to the survey.

Less than 1 in 20 students in surveyed SVVSD high schools (4.6%) reported they had used smokeless tobacco (e.g., chewing tobacco, snuff, or dip) while on school property in the 30 days prior to the survey. Overall, the prevalence of having used smokeless tobacco was higher among white non-Hispanic students (5.8%) than Hispanic students (3.1%); it was also higher among male students than female students (8.7% and 0.4%, respectively).

Among students in surveyed SVVSD high schools, 3.1% reported they had drunk at least 1 drink of alcohol while on school property during the 30 days before the survey. Overall, the prevalence of having drunk alcohol on school property was higher among male students than female students (4.5% and 1.7%, respectively).



Peer Use of Cigarettes, Alcohol and Marijuana

Grade	Peers smoked cigarettes*			Peers drank alcohol without parents' awareness*			Peers used marijuana*		
	Total	Female	Male	Total	Female	Male	Total	Female	Male
9th	49.5% (46.8-52.2)	52.8% (48.8-56.7)	46.2% (42.6-49.9)	63.3% (60.7-65.9)	70.1% (66.4-73.5)	56.7% (52.9-60.3)	49.9% (47.2-52.6)	51.1% (47.1-55.0)	48.7% (45.0-52.5)
10th	43.6% (40.9-46.2)	42.0% (38.2-45.9)	45.1% (41.3-48.9)	64.9% (62.3-67.4)	67.7% (64.1-71.1)	62.2% (58.4-65.9)	49.8% (47.2-52.3)	47.7% (43.9-51.5)	51.8% (48.1-55.4)
11th	59.8% (57.0-62.4)	67.0% (63.0-70.7)	53.1% (49.3-56.9)	81.8% (79.6-83.8)	86.6% (83.9-88.9)	77.4% (74.0-80.5)	66.6% (63.9-69.2)	71.6% (67.8-75.2)	62.0% (58.2-65.7)
12th	61.0% (58.0-64.0)	63.8% (59.5-67.9)	58.4% (53.7-62.9)	84.8% (82.3-87.0)	85.9% (82.7-88.6)	83.8% (79.8-87.1)	73.5% (70.7-76.2)	77.8% (73.8-81.3)	69.4% (65.0-73.5)
Race/Ethnicity									
White non-Hispanic	51.2% (49.5-52.9)	---	---	72.2% (70.7-73.7)	---	---	60.3% (58.6-62.0)	---	---
Hispanic	61.7% (58.7-64.7)	---	---	80.1% (77.6-82.3)	---	---	65.1% (62.2-67.9)	---	---
Other non-Hispanic	47.7% (43.2-52.3)	---	---	68.1% (63.8-72.0)	---	---	46.6% (42.1-51.2)	---	---
Total	53.5% (52.1-54.9)	56.4% (54.4-58.4)	50.7% (48.7-52.7)	73.8% (72.5-74.9)	77.6% (76.0-79.2)	70.1% (68.2-71.8)	60.0% (58.6-61.4)	62.0% (60.1-64.0)	58.1% (56.1-60.0)

* Four best friends in 12 months preceding survey
 --- Not available

Students in surveyed SVVSD high schools were asked to think of their 4 best friends and determine if any of them had smoked cigarettes in the 12 months prior to the survey. More than half (53.5%) reported that at least 1 of their 4 friends had smoked cigarettes during the past 12 months. Overall, the prevalence of having a friend who had smoked cigarettes was higher among Hispanic students (61.7%) than both white non-Hispanic and other non-Hispanic students (51.2% and 47.7%, respectively); it was also higher among female students (56.4%) than male students (50.7%).

Students in surveyed SVVSD high schools were asked to determine how many of their 4 best friends had tried beer, wine, or hard liquor without their parents' knowledge in the 12 months prior to the survey. More than 7 in 10 students (73.8%) reported that at least 1 of these 4 friends had tried alcohol without parental knowledge. Overall, the prevalence for this behavior was higher among Hispanic students (80.1%) than both white non-Hispanic and other non-Hispanic students (72.2% and 68.1%, respectively); it was also higher among female students (77.6%) than male students (70.1%).

Students in surveyed SVVSD high schools recalled their 4 best friends and were asked how many of them had used marijuana in the 12 months before the survey. Three in 5 students (60.0%) reported that at least 1 of those 4 friends had used marijuana in the previous year. Overall, the prevalence for having a friend who had used marijuana in the 12 months before the survey was higher among Hispanic students (65.1%) than white non-Hispanic students (60.3%), and both were higher than other non-Hispanic students (46.6%). The prevalence was also higher among female students than male students (62.0% and 58.1%, respectively).



Sexual Behavior - Sexual Intercourse and Sexual Activity

Grade	Ever had sexual intercourse			Currently sexually active*			Four or more sex partners during lifetime		
	Total	Female	Male	Total	Female	Male	Total	Female	Male
9th	21.1% (18.9-23.4)	17.6% (14.8-20.7)	24.6% (21.4-28.1)	8.1% (6.8-9.7)	8.9% (6.9-11.4)	7.4% (5.7-9.5)	3.6% (2.8-4.7)	1.9% (1.1-3.1)	5.5% (4.0-7.3)
10th	25.1% (22.7-27.8)	25.6% (22.2-29.3)	24.7% (21.1-28.7)	15.3% (13.2-17.7)	17.8% (14.8-21.2)	12.8% (10.0-16.3)	5.7% (4.4-7.4)	6.3% (4.4-8.8)	5.2% (3.4-7.8)
11th	41.3% (38.5-44.2)	43.1% (38.9-47.5)	39.5% (35.8-43.5)	29.0% (26.4-31.7)	32.6% (28.6-36.8)	25.7% (22.4-29.4)	13.2% (11.3-15.4)	11.9% (9.3-15.1)	14.4% (11.7-17.7)
12th	60.0% (56.4-63.4)	57.6% (52.4-62.6)	62.3% (57.4-67.0)	39.7% (36.3-43.3)	38.8% (33.8-44.0)	40.7% (36.0-45.5)	16.2% (13.7-19.1)	12.0% (8.7-16.3)	20.3% (16.5-24.6)
Race/Ethnicity									
White non-Hispanic	34.1% (32.3-35.8)	---	---	22.0% (20.5-23.6)	---	---	9.3% (8.2-10.6)	---	---
Hispanic	48.7% (45.5-51.8)	---	---	28.8% (25.9-31.9)	---	---	12.7% (10.7-15.0)	---	---
Other non-Hispanic	29.0% (25.0-33.4)	---	---	16.9% (13.7-20.6)	---	---	5.4% (3.6-8.3)	---	---
Total	37.0% (35.6-38.4)	35.9% (33.8-38.0)	38.0% (36.0-40.1)	23.1% (21.8-24.4)	24.3% (22.4-26.3)	21.8% (20.1-23.7)	9.7% (8.8-10.7)	8.0% (6.7-9.4)	11.5% (10.1-13.0)

* At least 1 sexual partner in 3 months preceding the survey

--- Not available

More than 1 in 3 students in surveyed SVVSD high schools (37.0%) reported they had ever had sexual intercourse. Overall, the prevalence of having had sexual intercourse was higher among Hispanic students (48.7%) than both white non-Hispanic (34.1%) and other non-Hispanic students (29.0%).

More than 1 in 5 students in surveyed SVVSD high schools (23.1%) reported they were currently sexually active (i.e., had sex with 1 or more partners within the 3 months preceding the survey). Overall, the prevalence of being sexually active was higher among Hispanic students (28.8%) than both white non-Hispanic (22.0%) and other non-Hispanic students (16.9%). Being sexually active increased with each grade level.

Nearly 1 in 10 students in surveyed SVVSD high schools (9.7%) reported they had had 4 or more sexual partners in their life. Overall, the prevalence of having 4 or more sexual partners was higher among Hispanic students (12.7%) than both white non-Hispanic and other non-Hispanic students (9.3% and 5.4%, respectively); it was also higher among male students (11.5%) than female students (8.0%).



Sexual Behavior - First Intercourse and Parental Perceptions

Grade	First sexual intercourse under age 13			First sexual partner was 5 or more years older or younger*			Parents feel it would be wrong for you to have sex		
	Total	Female	Male	Total	Female	Male	Total	Female	Male
9th	5.0% (3.9-6.4)	3.2% (1.9-5.3)	6.9% (5.2-9.0)	7.5% (4.8-11.5)	***	13.0% (8.4-19.7)	81.8% (79.7-83.7)	92.9% (90.9-94.5)	70.8% (67.3-74.0)
10th	2.1% (1.4-3.4)	***	3.5% (2.1-5.8)	6.6% (4.4-9.9)	***	8.9% (5.2-14.8)	72.0% (69.4-74.5)	82.8% (79.5-85.7)	61.6% (57.7-65.4)
11th	2.8% (2.0-4.1)	***	4.1% (2.7-6.3)	6.8% (4.9-9.3)	10.6% (7.2-15.4)	***	64.7% (61.9-67.4)	75.3% (71.5-78.7)	54.9% (51.0-58.7)
12th	3.9% (3.0-5.1)	2.1% (1.4-3.2)	5.6% (4.0-7.8)	5.2% (3.2-8.6)	7.6% (3.8-14.8)	3.1% (1.9-5.2)	53.8% (50.4-57.1)	58.4% (53.3-63.3)	49.3% (44.6-54.0)
Race/Ethnicity									
White non-Hispanic	2.3% (1.8-2.9)	---	---	5.6% (4.0-7.9)	---	---	68.7% (66.9-70.3)	---	---
Hispanic	6.7% (5.3-8.4)	---	---	7.6% (5.6-10.3)	---	---	63.7% (60.6-66.6)	---	---
Other non-Hispanic	3.7% (2.2-6.4)	---	---	***	---	---	75.3% (71.0-79.1)	---	---
Total	3.5% (3.0-4.1)	1.9% (1.4-2.5)	5.0% (4.2-6.0)	6.2% (5.0-7.8)	7.0% (4.9-9.9)	5.6% (4.3-7.1)	68.1% (66.7-69.5)	77.4% (75.5-79.2)	59.1% (57.1-61.1)

* Among the 37.0% of students who reported ever having had sexual intercourse

*** Numerator <3 and/or denominator <30

--- Not available

Among students in surveyed SVVSD high schools, 3.5% reported their first sexual intercourse was before the age of 13. Overall, prevalence of having had sex before age 13 was higher among Hispanic students (6.7%) than white non-Hispanic students (2.3%); it was also higher among male students (5.0%) than female students (1.9%).

Among students in surveyed SVVSD high schools, 6.2% reported that their first sexual partner was at least 5 years older or younger than them.

Nearly 7 in 10 students in surveyed SVVSD high schools (68.1%) reported that their parents/guardians felt it was wrong or very wrong for them to have sex. Overall, the prevalence of this belief was higher among other non-Hispanic students (75.3%) than white non-Hispanic students (68.7%), and both were higher than Hispanic students (63.7%). Prevalence was higher among female students (77.4%) than male students (59.1%). Students believe that their parents/guardians felt it was either wrong or very wrong for high school students to have sex decreased as grade level increased.



Sexual Behavior - Alcohol/Drugs and Condom Use

Grade	Alcohol or drugs used before last sexual intercourse*			Condom use during last sexual intercourse*		
	Total	Female	Male	Total	Female	Male
9th	20.6% (14.1-29.2)	24.4% (14.7-37.8)	*** ***	*** ***	*** ***	*** ***
10th	28.6% (21.5-36.9)	28.4% (19.5-39.5)	28.8% (18.5-41.9)	62.6% (54.2-70.2)	*** ***	*** ***
11th	18.2% (14.3-22.9)	16.2% (11.2-22.8)	20.5% (14.8-27.7)	53.5% (47.9-59.0)	44.0% (36.7-51.7)	64.6% (56.8-71.7)
12th	18.0% (14.4-22.4)	17.2% (12.3-23.7)	18.8% (13.8-25.0)	70.0% (64.6-75.0)	68.0% (59.7-75.4)	71.9% (64.6-78.3)
Race/Ethnicity						
White non-Hispanic	16.5% (13.7-19.7)	---	---	62.1% (57.9-66.1)	---	---
Hispanic	26.3% (21.0-32.4)	---	---	69.5% (63.2-75.1)	---	---
Other non-Hispanic	*** ***	---	---	*** ***	---	---
Total	20.0% (17.5-22.8)	19.6% (16.1-23.5)	20.5% (17.0-24.5)	64.0% (60.7-67.2)	57.2% (52.4-61.8)	71.4% (66.9-75.5)

* Among the 23.1% who reported being currently sexually active

*** Numerator <3 and/or denominator <30

--- Not available

Among the 23.1% of students in surveyed SVVSD high schools who reported being currently sexually active at the time of the survey, 1 in 5 (20.0%) reported they had either drunk alcohol or used drugs before the last time they had sex. Overall, the prevalence of drinking alcohol or using drugs before they last had sex was higher among Hispanic students than white non-Hispanic students (26.3% and 16.5%, respectively).

Among the 23.1% of students in surveyed SVVSD high schools who reported being sexually active at the time of the survey, more than 3 in 5 (64.0%) reported that they or their partner used a condom the last time they had sex. Overall, the prevalence of reported condom use (by the student or their partner) was higher among male students than female students (71.4% and 57.2%, respectively).



Sexual Behavior - Birth Control and Pregnancy

Grade	Birth control pills at last sexual intercourse*			Has been pregnant or gotten someone pregnant		
	Total	Female	Male	Total	Female	Male
9th	*** ***	*** ***	*** ***	1.0% (0.6-1.7)	*** ***	*** ***
10th	17.9% (12.9-24.2)	*** ***	*** ***	3.1% (2.1-4.5)	3.1% (2.0-4.9)	3.1% (1.7-5.5)
11th	36.5% (31.2-42.1)	47.7% (40.1-55.3)	23.0% (16.7-30.8)	2.4% (1.7-3.5)	1.5% (1.0-2.1)	3.4% (2.1-5.3)
12th	28.5% (23.4-34.1)	39.2% (30.6-48.4)	19.4% (14.8-25.1)	4.1% (3.0-5.5)	2.7% (1.8-4.1)	5.4% (3.7-8.0)
Race/Ethnicity						
White non-Hispanic	31.9% (27.9-36.2)	---	---	2.1% (1.7-2.7)	---	---
Hispanic	20.9% (16.2-26.7)	---	---	4.4% (3.3-5.9)	---	---
Other non-Hispanic	*** ***	---	---	*** ***	---	---
Total	27.5% (24.5-30.8)	35.1% (30.4-40.1)	19.5% (16.2-23.4)	2.6% (2.2-3.2)	2.0% (1.6-2.5)	3.3% (2.5-4.2)

* Among the 23.1% who reported as being currently sexually active
 *** Numerator <3 and/or denominator <30
 --- Not available

Among the 23.1% of students in surveyed SVVSD high schools who reported being sexually active at the time of the survey, more than 1 in 4 (27.5%) reported that they or their partner used birth control pills to prevent pregnancy the last time they had sexual intercourse. Overall, the prevalence of reported birth control use (by the student or their partner) was higher among white non-Hispanic students (31.9%) than Hispanic students (20.9%); it was also higher among female students (35.1%) than male students (19.5%).

Among students in surveyed SVVSD high schools, 2.6% reported they had either been pregnant or gotten someone pregnant. Overall, the prevalence of having been pregnant or gotten someone pregnant was higher among Hispanic students (4.4%) than white non-Hispanic students (2.1%).





Body Weight - Overweight

	<i>Overweight*</i>			<i>Obese*</i>			<i>Thought they were overweight</i>		
Grade	Total	Female	Male	Total	Female	Male	Total	Female	Male
9th	9.2% (7.9-10.7)	9.1% (7.3-11.4)	9.2% (7.4-11.5)	7.3% (5.8-9.0)	5.8% (4.0-8.2)	8.7% (6.6-11.4)	24.0% (21.8-26.3)	28.5% (25.1-32.2)	19.5% (16.8-22.6)
10th	11.5% (9.9-13.3)	10.9% (8.7-13.6)	12.1% (9.9-14.7)	9.9% (8.2-11.9)	5.0% (3.4-7.2)	14.5% (11.7-17.8)	25.8% (23.4-28.3)	28.7% (25.4-32.3)	22.9% (19.7-26.5)
11th	11.2% (9.3-13.4)	8.7% (6.5-11.5)	13.6% (10.7-17.1)	6.4% (5.0-8.2)	3.2% (2.0-5.3)	9.4% (7.1-12.4)	25.2% (22.8-27.7)	27.8% (24.2-31.7)	22.7% (19.8-26.0)
12th	12.3% (10.4-14.5)	6.0% (4.4-8.0)	19.1% (15.6-23.2)	9.4% (7.5-11.6)	4.9% (3.1-7.8)	14.2% (11.1-17.9)	31.1% (28.3-34.0)	33.2% (29.2-37.5)	29.1% (25.1-33.3)
Race/Ethnicity									
White non-Hispanic	10.7% (9.6-11.9)	---	---	5.3% (4.5-6.1)	---	---	24.9% (23.4-26.4)	---	---
Hispanic	12.7% (10.9-14.8)	---	---	16.2% (13.8-18.9)	---	---	30.9% (28.1-33.9)	---	---
Other non-Hispanic	9.8% (7.6-12.7)	---	---	9.3% (6.5-13.1)	---	---	26.6% (22.8-30.9)	---	---
Total	11.0% (10.2-12.0)	8.6% (7.6-9.8)	13.4% (12.0-14.9)	8.2% (7.4-9.2)	4.7% (3.8-5.8)	11.6% (10.3-13.2)	26.5% (25.3-27.8)	29.5% (27.7-31.4)	23.6% (21.9-25.3)

* Overweight = Body Mass Index (BMI) at or above the 85th percentile and below the 95th percentile; Obese = BMI at or above the 95th percentile
 --- Not available

More than 1 in 10 students in surveyed SVVSD high schools (11.0%) were categorized as being overweight, based on their self-reported height and weight (without shoes). Overall, the prevalence of being overweight was higher among male students (13.4%) than female students (8.6%).

Among students in surveyed SVVSD high schools, 8.2% were categorized as being obese, based upon the height and weight they reported in the survey. Overall, the prevalence of obesity was higher among Hispanic students (16.2%) than other non-Hispanic students (9.3%), and both were higher than white non-Hispanic students (5.3%). The prevalence of obesity was higher among male students (11.6%) than female students (4.7%).

When asked to describe their weight, more than 1 in 4 students in surveyed SVVSD high schools (26.5%) reported they thought of themselves as being overweight. Overall, the prevalence of that belief was higher among Hispanic students (30.9%) than white non-Hispanic students (24.9%); it was also higher among female students than male students (29.5% and 23.6%, respectively).



Body Weight - Weight Loss

Grade	Attempting weight loss*			Exercised to lose weight or avoid gaining weight*			Ate less food, fewer calories, or foods low in fat to avoid weight gain*		
	Total	Female	Male	Total	Female	Male	Total	Female	Male
9th	40.3% (37.7-42.9)	52.2% (48.2-56.2)	28.5% (25.3-32.0)	59.4% (56.7-62.1)	74.5% (70.8-77.9)	44.5% (41.0-48.2)	32.4% (29.8-35.0)	46.8% (42.9-50.8)	18.2% (15.5-21.3)
10th	42.9% (40.1-45.7)	53.6% (49.7-57.5)	32.4% (28.7-36.3)	59.1% (56.5-61.7)	70.6% (67.0-73.9)	47.9% (44.1-51.7)	33.9% (31.4-36.6)	45.6% (41.8-49.5)	22.6% (19.4-26.1)
11th	42.6% (39.7-45.5)	57.2% (52.9-61.3)	29.3% (25.8-33.1)	58.1% (55.2-60.9)	72.1% (68.2-75.7)	45.2% (41.2-49.1)	37.7% (35.0-40.5)	48.6% (44.4-52.8)	27.7% (24.2-31.4)
12th	47.4% (43.9-50.8)	60.0% (55.5-64.4)	35.0% (30.5-39.8)	61.7% (58.3-65.0)	68.7% (64.3-72.8)	55.0% (50.2-59.7)	41.6% (38.3-44.9)	53.8% (49.3-58.3)	29.8% (25.8-34.2)
Race/Ethnicity									
White non-Hispanic	39.4% (37.7-41.2)	---	---	56.5% (54.7-58.3)	---	---	35.3% (33.6-37.0)	---	---
Hispanic	53.7% (50.5-56.9)	---	---	69.1% (66.1-71.9)	---	---	42.4% (39.4-45.5)	---	---
Other non-Hispanic	44.4% (39.9-49.0)	---	---	58.7% (54.3-63.1)	---	---	31.1% (27.1-35.4)	---	---
Total	43.2% (41.8-44.7)	55.7% (53.6-57.8)	31.3% (29.4-33.2)	59.6% (58.1-61.0)	71.5% (69.6-73.3)	48.1% (46.1-50.1)	36.4% (35.0-37.8)	48.7% (46.6-50.8)	24.6% (22.9-26.4)

* During the 30 days preceding the survey
 --- Not available

More than 2 in 5 students in surveyed SVVSD high schools (43.2%) reported they had tried to lose weight in the 30 days preceding the survey. Overall, the prevalence of attempted weight loss was higher among Hispanic students (53.7%) than both other non-Hispanic and white non-Hispanic students (44.4% and 39.4%, respectively). Additionally, at each of the four grade levels, female students (55.7%) had a higher prevalence of having attempted weight loss than male students (31.3%).

Nearly 3 in 5 students in surveyed SVVSD high schools (59.6%) reported they had exercised to lose weight or to keep from gaining weight in the 30 days prior to the survey. Overall, the prevalence of having exercised for weight loss was higher among Hispanic students (69.1%) than both other non-Hispanic and white non-Hispanic students (58.7% and 56.5%, respectively). Additionally, at each of the four grade levels, female students (48.7%) had a higher prevalence of exercising for weight loss than male students (24.6%).

Approximately 1 in 3 students in surveyed SVVSD high schools (36.4%) reported they had eaten less food, fewer calories, or foods low in fat to lose weight or to keep from gaining weight in the 30 days before the survey. Overall, the prevalence of dietary changes to lose weight was higher among Hispanic students (42.4%) than both white non-Hispanic and other non-Hispanic students (35.3% and 31.1%, respectively). Additionally, at each of the four grade levels, female students (48.7%) had a higher prevalence of dietary changes to lose weight than male students (24.6%).



Body Weight - Unhealthy Weight Loss Practices

Grade	<i>Fasting 24 hours or more to lose weight or to avoid gaining weight*</i>			<i>Took diet pills, powders, or liquids to lose weight or to avoid gaining weight**</i>			<i>Vomited or took laxatives to lose weight or to avoid gaining weight*</i>		
	Total	Female	Male	Total	Female	Male	Total	Female	Male
9th	8.5% (7.1-10.2)	12.4% (10.0-15.2)	4.7% (3.3-6.8)	1.0% (0.6-1.7)	*** ***	*** ***	1.6% (1.1-2.3)	2.6% (1.7-3.9)	*** ***
10th	8.6% (7.1-10.5)	13.2% (10.6-16.2)	4.2% (2.7-6.5)	3.8% (2.7-5.3)	5.9% (4.0-8.7)	1.8% (1.1-2.8)	4.8% (3.6-6.4)	8.2% (6.0-11.2)	*** ***
11th	9.3% (7.8-11.2)	12.9% (10.4-15.9)	6.1% (4.3-8.5)	5.6% (4.4-7.1)	8.0% (6.0-10.7)	3.3% (2.1-5.2)	4.9% (4.0-6.1)	6.9% (5.3-9.0)	3.1% (2.2-4.5)
12th	10.3% (8.4-12.4)	12.2% (9.4-15.7)	8.4% (6.3-11.1)	2.7% (2.1-3.6)	2.2% (1.5-3.2)	3.2% (2.2-4.8)	2.9% (2.0-4.2)	3.4% (2.3-4.9)	2.5% (1.2-4.8)
Race/Ethnicity									
White non-Hispanic	8.1% (7.1-9.1)	---	---	2.8% (2.3-3.4)	---	---	3.4% (2.9-4.1)	---	---
Hispanic	13.5% (11.5-15.8)	---	---	5.1% (3.8-6.8)	---	---	5.1% (3.8-6.7)	---	---
Other non-Hispanic	6.3% (4.6-8.7)	---	---	2.4% (1.4-4.0)	---	---	*** ***	---	---
Total	9.2% (8.3-10.1)	12.7% (11.3-14.2)	5.9% (4.9-7.0)	3.3% (2.8-3.8)	4.4% (3.6-5.3)	2.3% (1.8-2.9)	3.6% (3.1-4.1)	5.3% (4.5-6.2)	1.9% (1.4-2.6)

* During the 30 days preceding the survey

** During the 30 days preceding the survey (not including meal replacement products such as Slim Fast)

*** Numerator <3 and/or denominator <30

--- Not available

Among students in surveyed SVVSD high schools, 9.2% reported that, in the 30 days before the survey, they had gone 24 hours or more without eating (i.e., fasting) in an effort to lose or maintain weight. Overall, the prevalence of having fasted for weight loss was higher among Hispanic students (13.5%) than both white non-Hispanic and other non-Hispanic students (8.1% and 6.3%, respectively); it was also higher among female students (12.7%) than male students (5.9%).

Among students in surveyed SVVSD high schools, 3.3% reported they had taken diet pills, powders, or liquids without a doctor's advice in an effort to lose weight or maintain weight in the 30 days prior to the survey. Overall, the prevalence of that behavior was higher among Hispanic students (5.1%) than white non-Hispanic students (2.8%); it was also higher among female students (4.4%) than male students (2.3%).

Nearly 1 in 25 students in surveyed SVVSD high schools (3.6%) reported they had vomited or taken laxatives to lose or maintain weight in the 30 days prior to the survey. Overall, the prevalence of that behavior was higher among female students than male students (5.3% and 1.9%, respectively).



Physical Activity and Sleep

Grade	Participated in vigorous physical activity*			Sleeps 8 or more hours on a school night		
	Total	Female	Male	Total	Female	Male
9th	75.5% (73.0-77.8)	67.6% (63.7-71.2)	83.3% (80.1-86.0)	52.5% (49.7-55.2)	37.8% (33.9-41.9)	67.2% (63.7-70.5)
10th	75.4% (72.8-77.7)	73.2% (69.5-76.7)	77.4% (74.0-80.6)	40.8% (38.1-43.5)	32.5% (29.0-36.3)	48.9% (45.1-52.8)
11th	72.7% (70.0-75.2)	63.8% (59.7-67.7)	81.0% (77.5-84.0)	34.1% (31.4-36.9)	34.2% (30.3-38.3)	34.0% (30.3-37.9)
12th	71.4% (68.0-74.6)	60.7% (55.5-65.5)	81.9% (77.4-85.6)	32.7% (29.6-36.0)	22.2% (18.7-26.1)	43.0% (38.3-47.9)
Race/Ethnicity						
White non-Hispanic	76.8% (75.1-78.4)	---	---	40.8% (39.0-42.5)	---	---
Hispanic	66.5% (63.5-69.4)	---	---	43.4% (40.4-46.6)	---	---
Other non-Hispanic	72.2% (67.9-76.2)	---	---	28.0% (24.2-32.0)	---	---
Total	73.7% (72.3-75.1)	66.3% (64.2-68.3)	80.9% (79.1-82.6)	39.9% (38.5-41.4)	31.7% (29.8-33.7)	48.0% (46.0-50.0)

* For at least 60 minutes on 3 or more of the 7 days preceding the survey.
 --- Not available

More than 7 in 10 students in surveyed SVVSD high schools (73.7%) reported they had participated in at least 60 minutes of vigorous physical activity on 3 or more of the 7 days preceding the survey. Overall, the prevalence of participating in vigorous physical activity was higher among white non-Hispanic students (76.8%) than Hispanic students (66.5%); it was also higher among male students (80.9%) than female students (66.3%).

Almost 2 in 5 students in surveyed SVVSD high schools (39.9%) reported they got 8 or more hours of sleep on an average school night. Overall, the prevalence of getting 8 or more hours of sleep on an average school night was higher among Hispanic and white non-Hispanic students (43.4% and 40.8%, respectively) than other non-Hispanic students (28.0%); it was also higher among male students than female students (48.0% and 31.7%, respectively).



Screen Time

	<i>3 or more hours of TV per day*</i>			<i>3 or more hours of video/computer games per day*</i>		
Grade	Total	Female	Male	Total	Female	Male
9th	23.9% (21.6-26.3)	22.5% (19.2-26.1)	25.3% (22.2-28.6)	23.7% (21.5-26.1)	23.4% (20.0-27.2)	24.0% (21.2-27.1)
10th	25.7% (23.3-28.3)	24.0% (20.8-27.4)	27.5% (24.0-31.4)	25.2% (22.8-27.7)	16.6% (13.9-19.7)	33.4% (29.9-37.3)
11th	22.9% (20.5-25.6)	23.9% (20.1-28.1)	22.0% (18.9-25.5)	23.8% (21.4-26.3)	17.6% (14.8-20.9)	29.6% (26.1-33.3)
12th	25.5% (22.6-28.6)	21.9% (18.4-25.8)	29.0% (24.7-33.8)	23.6% (20.7-26.8)	16.3% (12.4-21.0)	30.7% (26.7-35.1)
Race/Ethnicity						
White non-Hispanic	19.6% (18.2-21.2)	---	---	24.4% (22.9-26.0)	---	---
Hispanic	36.3% (33.3-39.3)	---	---	22.8% (20.1-25.6)	---	---
Other non-Hispanic	26.9% (22.8-31.4)	---	---	24.2% (20.6-28.2)	---	---
Total	24.5% (23.2-25.8)	23.0% (21.3-24.9)	25.9% (24.1-27.8)	24.1% (22.8-25.4)	18.5% (16.8-20.3)	29.4% (27.6-31.3)

* On an average school day

--- Not available

Nearly 1 in 4 students in surveyed SVVSD high schools (24.5%) reported they watched 3 or more hours of TV on an average school day. Overall, the prevalence of watching 3 or more hours of TV per day was higher among Hispanic students (36.3%) than other non-Hispanic students (26.9%), and both were higher than white non-Hispanic students (19.6%).

Nearly 1 in 4 students in surveyed SVVSD high schools (24.1%) reported they played video or computer games or used a computer for something other than school work for 3 or more hours on an average school day. Overall, the prevalence was higher among male students (29.4%) than female students (18.5%).

APPENDIX 1

YRBS 2009 STATE AND NATIONAL DATA

		Total	Female	Male
Parents had clear rules and standards for behavior*	Colorado	NA	NA	NA
	National	NA	NA	NA
Ate 4 or more meals with family*	Colorado	NA	NA	NA
	National	NA	NA	NA
Did community service*	Colorado	NA	NA	NA
	National	NA	NA	NA
Rarely or never wore bicycle helmet	Colorado	75.2%	71.9%	77.7%
	National	84.7%	83.1%	85.8%
Rarely or never wore seatbelt	Colorado	7.6%	5.5%	9.7%
	National	9.7%	7.7%	11.5%
Rode with a driver who had been drinking alcohol	Colorado	24.6%	25.9%	23.4%
	National	28.3%	28.8%	27.8%
Drove after drinking alcohol	Colorado	7.4%	7.0%	7.8%
	National	9.7%	7.6%	11.6%
Felt safe in their neighborhood*	Colorado	NA	NA	NA
	National	NA	NA	NA
Felt too unsafe to go to school	Colorado	5.1%	4.8%	5.5%
	National	5.0%	5.3%	4.6%
Harassed*	Colorado	NA	NA	NA
	National	NA	NA	NA
Harassed because of race or ethnic origin*	Colorado	NA	NA	NA
	National	NA	NA	NA
Harassed because someone thought they were lesbian, gay, or bisexual*	Colorado	NA	NA	NA
	National	NA	NA	NA
Harassed because of religious beliefs*	Colorado	NA	NA	NA
	National	NA	NA	NA
Harassed because of weight, size, physical appearance*	Colorado	NA	NA	NA
	National	NA	NA	NA
Received unwanted sexual comments or attention*	Colorado	NA	NA	NA
	National	NA	NA	NA
Carried a weapon	Colorado	16.7%	8.2%	25.0%
	National	17.5%	7.1%	27.1%
In a physical fight	Colorado	32.0%	21.8%	42.0%
	National	31.5%	22.9%	39.3%

* Survey question was used only by Boulder County

APPENDIX 1 (CONT'D)

YRBS 2009 STATE AND NATIONAL DATA

		Total	Female	Male
Injured in fight and needed medical treatment	Colorado	4.0%	3.3%	4.7%
	National	3.8%	2.2%	5.1%
Carried a weapon on school property	Colorado	5.5%	3.0%	8.0%
	National	5.6%	2.9%	8.0%
Threatened or injured with a weapon on school property	Colorado	8.0%	5.1%	10.8%
	National	7.7%	5.5%	9.6%
In physical fight on school property	Colorado	10.7%	5.0%	16.2%
	National	11.1%	6.7%	15.1%
Hurt by boyfriend/girlfriend	Colorado	9.1%	6.9%	11.2%
	National	9.8%	9.3%	10.3%
Electronically bullied*	Colorado	NA	NA	NA
	National	NA	NA	NA
Ever forced to have sexual intercourse	Colorado	7.7%	11.1%	4.4%
	National	7.4%	10.5%	4.5%
Ever touched sexually when did not want to be touched*	Colorado	NA	NA	NA
	National	NA	NA	NA
Felt sad or hopeless	Colorado	25.4%	31.7%	19.3%
	National	26.1%	33.9%	19.1%
Had someone to talk to when feeling sad or hopeless*	Colorado	NA	NA	NA
	National	NA	NA	NA
Usually got help when feeling sad or hopeless*	Colorado	NA	NA	NA
	National	NA	NA	NA
Intentionally self-injured*	Colorado	NA	NA	NA
	National	NA	NA	NA
Seriously considered attempting suicide	Colorado	13.7%	17.9%	9.7%
	National	13.8%	17.4%	10.5%
Planned suicide	Colorado	11.0%	12.8%	9.3%
	National	10.9%	13.2%	8.6%
Attempted suicide	Colorado	7.6%	9.3%	5.6%
	National	6.3%	8.1%	4.6%
Suicide attempt required medical treatment	Colorado	3.1%	3.5%	2.4%
	National	1.9%	2.3%	1.6%
Ever smoked cigarettes	Colorado	43.1%	41.4%	44.6%
	National	46.3%	46.1%	46.3%

* Survey question was used only by Boulder County

APPENDIX 1 (CONT'D)

YRBS 2009 STATE AND NATIONAL DATA

		Total	Female	Male
Current cigarette user	Colorado	17.7%	17.4%	17.8%
	National	19.5%	19.1%	19.8%
Current frequent cigarette use	Colorado	10.7%	6.2%	15.0%
	National	7.8%	4.1%	11.1%
Bought cigarettes in a store or gas station	Colorado	12.6%	10.3%	**
	National	14.1%	9.6%	18.3%
Ever smoked cigarettes daily	Colorado	10.6%	10.7%	10.5%
	National	11.2%	10.6%	11.7%
Tried to quit smoking cigarettes	Colorado	53.2%	60.6%	46.4%
	National	50.8%	54.2%	48.0%
Current smokeless tobacco use	Colorado	10.7%	4.2%	16.8%
	National	8.9%	2.2%	15.0%
Current cigar use	Colorado	16.3%	11.4%	21.0%
	National	14.0%	8.8%	18.6%
Current tobacco use	Colorado	26.5%	21.6%	31.3%
	National	26.0%	21.8%	29.8%
Felt smoking 1+ packs per day was moderate or great risk*	Colorado	NA	NA	NA
	National	NA	NA	NA
Felt it was wrong or very wrong for someone their age to smoke*	Colorado	NA	NA	NA
	National	NA	NA	NA
Parents felt it was wrong or very wrong for you to smoke*	Colorado	NA	NA	NA
	National	NA	NA	NA
Ever drank alcohol	Colorado	72.4%	77.3%	67.8%
	National	72.5%	74.2%	70.8%
Current alcohol use	Colorado	40.8%	44.2%	37.4%
	National	41.8%	42.9%	40.8%
Binge drinking	Colorado	25.1%	27.2%	23.2%
	National	24.2%	23.4%	25.0%
Felt drinking 1-2 drinks per day was moderate or great risk*	Colorado	NA	NA	NA
	National	NA	NA	NA
Felt it was wrong or very wrong for someone their age to drink*	Colorado	NA	NA	NA
	National	NA	NA	NA
Parents felt it was wrong or very wrong for you to drink*	Colorado	NA	NA	NA
	National	NA	NA	NA

* Survey question was used only by Boulder County

** Less than 100 students in the subgroup

APPENDIX 1 (CONT'D)

YRBS 2009 STATE AND NATIONAL DATA

		Total	Female	Male
Ever used marijuana	Colorado	42.6%	40.8%	44.5%
	National	36.8%	34.3%	39.0%
Current marijuana use	Colorado	24.8%	24.2%	25.5%
	National	20.8%	17.9%	23.4%
Felt smoking marijuana regularly was a moderate or great risk*	Colorado	NA	NA	NA
	National	NA	NA	NA
Felt it was wrong or very wrong for someone their age to smoke marijuana*	Colorado	NA	NA	NA
	National	NA	NA	NA
Parents feel it is wrong or very wrong for you to smoke marijuana*	Colorado	NA	NA	NA
	National	NA	NA	NA
Smoked a whole cigarette before age 13	Colorado	8.8%	7.5%	10.0%
	National	10.7%	9.4%	11.8%
Drank alcohol before age 13	Colorado	22.4%	17.5%	27.2%
	National	21.1%	18.1%	23.7%
Tried marijuana before age 13	Colorado	8.3%	6.6%	10.0%
	National	7.5%	5.0%	9.7%
Offered, sold, or was given an illegal drug on school property	Colorado	22.7%	19.7%	25.7%
	National	22.7%	19.3%	25.9%
Attended school under the influence of alcohol or other illegal drugs*	Colorado	NA	NA	NA
	National	NA	NA	NA
Smoked cigarettes on school property	Colorado	8.3%	7.7%	8.8%
	National	5.1%	4.0%	6.2%
Used smokeless tobacco on school property	Colorado	5.9%	2.6%	9.1%
	National	5.5%	1.1%	9.4%
Drank alcohol on school property	Colorado	4.1%	3.8%	4.5%
	National	4.5%	3.6%	5.3%
Peers smoked cigarettes*	Colorado	NA	NA	NA
	National	NA	NA	NA
Peers drank alcohol without parents' awareness*	Colorado	NA	NA	NA
	National	NA	NA	NA
Peers used marijuana*	Colorado	NA	NA	NA
	National	NA	NA	NA
Ever used cocaine	Colorado	8.3%	7.2%	9.4%
	National	6.4%	5.3%	7.3%

* Survey question was used only by Boulder County

APPENDIX 1 (CONT'D)

YRBS 2009 STATE AND NATIONAL DATA

		Total	Female	Male
Ever used inhalants	Colorado	9.2%	9.9%	8.6%
	National	11.7%	12.9%	10.6%
Ever used ecstasy	Colorado	10.2%	9.1%	11.4%
	National	6.7%	5.5%	7.6%
Ever used heroin	Colorado	3.2%	2.1%	4.1%
	National	2.5%	1.7%	3.2%
Ever used methamphetamines	Colorado	3.5%	2.8%	4.1%
	National	4.1%	3.3%	4.7%
Ever used a needle to inject drugs	Colorado	2.7%	2.1%	3.1%
	National	2.1%	1.4%	2.7%
Ever used illegal steroids	Colorado	3.6%	2.9%	4.3%
	National	3.3%	2.2%	4.3%
Ever used prescription drugs without a prescription	Colorado	NA	NA	NA
	National	20.2%	19.8%	20.4%
Ever used other illegal drugs*	Colorado	NA	NA	NA
	National	NA	NA	NA
Current cocaine use	Colorado	3.6%	3.3%	4.0%
	National	2.8%	2.0%	3.5%
Ever had sexual intercourse	Colorado	40.0%	36.2%	43.7%
	National	46.0%	45.7%	46.1%
Currently sexually active	Colorado	27.4%	25.6%	29.1%
	National	34.2%	35.6%	32.6%
Four or more sex partners during lifetime	Colorado	13.8%	11.0%	16.5%
	National	13.8%	11.2%	16.2%
First sexual intercourse under age 13	Colorado	4.6%	2.7%	6.4%
	National	5.9%	3.1%	8.4%
First sexual partner was 5 or more years older/younger*	Colorado	NA	NA	NA
	National	NA	NA	NA
Parents feel it would be wrong for you to have sex	Colorado	NA	NA	NA
	National	NA	NA	NA
Alcohol or drug use at last sexual intercourse	Colorado	24.4%	26.7%	22.6%
	National	21.6%	17.1%	25.9%
Condom use during last sexual intercourse	Colorado	63.2%	59.0%	66.6%
	National	61.1%	53.9%	68.6%

* Survey question was used only by Boulder County

APPENDIX 1 (CONT'D)

YRBS 2009 STATE AND NATIONAL DATA

		Total	Female	Male
Used birth control pills at last sexual intercourse	Colorado	22.3%	25.6%	19.4%
	National	19.8%	23.0%	16.5%
Have been pregnant or gotten someone pregnant*	Colorado	NA	NA	NA
	National	NA	NA	NA
Overweight	Colorado	11.1%	10.4%	11.7%
	National	15.8%	15.9%	15.7%
Obese	Colorado	7.1%	5.4%	8.6%
	National	12.0%	8.3%	15.3%
Thought they were overweight	Colorado	21.9%	26.1%	18.0%
	National	27.7%	33.1%	22.7%
Attempting weight loss	Colorado	38.6%	52.8%	24.8%
	National	44.4%	59.3%	30.5%
Exercised to lose weight or avoid gaining weight	Colorado	58.6%	68.7%	49.2%
	National	61.5%	67.9%	55.7%
Ate less food, fewer calories, or foods low in fat to avoid weight gain	Colorado	32.4%	43.6%	21.5%
	National	39.5%	51.6%	28.4%
Fasted 24 hours or more to lose weight or avoid gaining weight	Colorado	9.8%	14.5%	5.3%
	National	10.6%	14.5%	6.9%
Took diet pills, powders, or liquids to lose weight or avoid gaining weight	Colorado	4.7%	5.8%	3.4%
	National	5.0%	6.3%	3.8%
Vomited or took laxatives to lose weight or avoid gaining weight	Colorado	3.6%	4.6%	2.6%
	National	4.0%	5.4%	2.6%
Participated in vigorous physical activity*	Colorado	NA	NA	NA
	National	NA	NA	NA
Slept 8 or more hours on school nights	Colorado	NA	NA	NA
	National	30.9%	28.2%	33.3%
Watched 3 or more hours of TV per day	Colorado	25.1%	20.5%	29.7%
	National	32.8%	32.1%	33.5%
Watched 3 or more hours of video/computer games per day	Colorado	18.4%	13.2%	23.3%
	National	24.9%	21.2%	28.3%

* Survey question was used only by Boulder County